

Aluminium in an ocean general circulation model compared with the West Atlantic Geotraces cruises*

M.M.P. van Hulten¹, A. Sterl¹, A. Tagliabue², J.-C. Dutay³, M. Gehlen³, H.J.W.
de Baar^{4,5}, R. Middag^{5,6}

¹Royal Netherlands Meteorological Institute (KNMI), ²University of
Cape Town (UCT), ³Laboratoire des Sciences du Climat et de
l'Environnement (LSCE), ⁴University of Groningen (RUG), ⁵Royal
Netherlands Institute for Sea Research (NIOZ), ⁶University of
California Santa Cruz (UCSC)

February 22, 2012

*This work is licensed under *Creative Commons Attribution-ShareAlike 3.0 Unported License*
(CC BY-SA). This note overrides other licenses, giving you true freedom for this preprint.

Abstract

A model of aluminium has been developed and implemented in an Ocean General Circulation Model (NEMO-PISCES). In the model, aluminium enters the ocean by means of dust deposition. The internal oceanic processes are described by advection, mixing and reversible scavenging. The model has been evaluated against a number of selected high-quality datasets covering much of the world ocean, especially those from the West Atlantic Geotraces cruises of 2010 and 2011. Generally, the model results are in fair agreement with the observations. However, the model does not describe well the vertical distribution of dissolved Al in the North Atlantic Ocean. The model may require changes in the physical forcing and the vertical dependency of the sinking velocity of biogenic silica to account for other discrepancies. To explore the model behaviour, sensitivity experiments have been performed, in which we changed the key parameters of the scavenging process as well as the input of aluminium into the ocean. This resulted in a better understanding of aluminium in the ocean, and it is now clear which parameter has what effect on the dissolved aluminium distribution and which processes might be missing in the model, among which boundary scavenging and biological incorporation of aluminium into diatoms.

Keywords: Aluminium, Dust deposition, Modelling, PISCES, Biogenic silica, GEOTRACES

NEMO Nucleus for European Modelling of the Ocean
OPA Océan PARallélisé
BEC Biogeochemical Elemental Cycling
INCA Integrated Nitrogen model for multiple source assessment in CAtchments
LMDzT Laboratoire de Météorologie Dynamique GCM
HAMOCC HAmburg Model for the Ocean Carbon Cycle
LSG (Hamburg) Large Scale Geostrophic model
OGCM Ocean General Circulation Model
IPY International Polar Year
BATS Bermuda Atlantic Time-series Study
SAFe Sampling and Analysis of Fe
POM Particulate Organic Matter
NADW North Atlantic Deep Water
AABW AntArctic Bottom Water
AAIW AntArctic Intermediate Water
SAAMW SubAntArctic Mode Water
MOC Meridional Overturning Circulation

1 Introduction

The distribution and cycling of aluminium (Al) in the ocean has received attention for a variety of reasons. Firstly, if the Al cycle is understood well, aluminium surface concentrations can be used to constrain atmospheric dust deposition fields (Gehlen et al. (2003); Measures et al. (2005); Han et al. (2008); Measures et al. (2010); Han (2010)), which are used to predict aeolian iron addition to the euphotic zone. This is important, since iron is an essential trace-nutrient for phytoplankton, thus its availability has a direct consequence on primary production and air-sea CO₂ exchange. Secondly, there is evidence that Al inhibits the solubility of sedimentary biogenic silica (Lewin (1961); Bennekom et al. (1991); Dixit et al. (2001); Sarmiento and Gruber (2006); Emerson and Hedges (2006)). If less biogenic silica gets dissolved from sediments, eventually there will be less silicic acid available in the euphotic zone, which will reduce diatom production as silicon is an essential major nutrient for diatoms. Modified diatom productivity will impact ocean food webs and the export of organic carbon to the ocean's interior. For advancement in both of these fields of interest a good understanding of the Al cycle is pertinent.

Currently it is assumed that the major source of Al to the ocean is via dust deposition (e.g. Orians and Bruland (1986); Maring and Duce (1987); Kramer et al. (2004); Measures et al. (2005)). When dust enters the ocean, a part of its aluminium content (1–15%) dissolves in the uppermost layer and is quickly distributed over the mixed layer by turbulent mixing. Most dust remains in the particulate phase and sinks to the bottom of the ocean, while a small fraction might dissolve in the water column. The dust that does not dissolve at all is buried in the sediment (Gehlen et al. (2003); Sarmiento and Gruber (2006)).

Arguments that dissolution occurs primarily in the upper layer of the ocean come from shipboard experiments and atmospheric moisture considerations. Maring and Duce (1987) and Measures et al. (2010) showed that within a day after deposition, most of dissolvable Al will be dissolved. Assuming a sinking speed of dust of 30 m/day, most Al would then dissolve in the upper 30 m of the ocean. This depth is shallower than the mixed layer depth, which means that there is little dissolution below the mixed layer. Even though some earlier studies showed that most deposition is dry (Jickells et al. (1994); Jickells (1995)), more recent work shows that dust deposition is mostly wet (Guerzoni et al. (1997); Vink and Measures (2001)). It has been argued that most dust passes through low *pH* environments in the atmosphere, which means that for wet deposition Al is already dissolved when it enters the ocean surface. Since the wetly deposited Al is most important, dissolution in the surface ocean is most relevant (as a lower bound, since dry deposition results in both surface and water column dissolution). Furthermore, based on the same low *pH* argument, dissolvable Al from dry deposition is likely to instantaneously dissolve in the surface ocean (Measures et al. (2010) and references herein), making the relative amount of surface dissolution even higher compared to water column dissolution.

Fluvial input can be thought to be important as well, since rivers carry large concentrations of Al, but in estuaries and coastal regions this Al is removed by scavenging of Al onto particles (Mackin (1986); Mackin and Aller (1986); Orians and Bruland (1986); Brown et al. (2010)). There are also indications for Al input as a consequence of sediment remobilisation, as in the Arctic Ocean (Middag et al. (2009)) and North Atlantic Ocean (Moran and Moore (1991)). However, the importance of sedimentary sources can vary by basin (e.g. in the Southern Ocean these are small as shown by Moran et al. (1992) and Middag et al. (2011)). Finally, hydrothermal vents are thought to only play a minor role (e.g. Hydes et al. (1986); Middag (2010)). In summary, the dominant external source of aluminium in the ocean is atmospheric dust deposition.

Dissolved aluminium (Al_{diss}) is removed mainly by particle scavenging (Orians and Bruland (1986); Moran and Moore (1989); Bruland and Lohan (2006)). This is the combination of adsorption onto a solid surface, followed by sinking due to insufficient buoyancy of the particulates in the seawater (Goldberg (1954); Bacon and Anderson (1982); Bruland and Lohan (2006)). Typically, scavenging is deemed to be reversible, which means that during sinking release of the adsorbed, or particulate, aluminium (Al_{ads}) may occur. This happens both directly (by desorption) and indirectly (by dissolution of the biogenic carrier particles). As a consequence, Al_{diss} concentrations increase with depth (Bacon and Anderson (1982); Anderson (2006)). In this way Al_{diss} is distributed over depth more efficiently than due to mixing and water mass transport. Aluminium is scavenged relatively efficiently and therefore has a relatively short residence time in the ocean (100–200 yr) (Orians and Bruland (1985)).

Except for scavenging, there are strong suggestions from observations in certain regions that Al is biologically incorporated into the siliceous cell walls of diatoms (Stoffyn (1979); Moran and Moore (1988); Gehlen et al. (2002)). It seems that Al does not play an essential role for the diatoms, but it can be incorporated functioning as a replacement for silicon (Si), since it is similar in size. Therefore it is likely that the incorporation ratio Al:Si is close to that of the surrounding waters. These regions include the Arctic Ocean (Middag et al. (2009)) and the Mediterranean Sea (MacKenzie et al. (1978); Hydes et al. (1988); Chou and Wollast (1997)). Given the ratio Al:Si of incorporation into the diatom in the photic zone, after remineralisation anywhere in the water column, the same dissolved Al:Si will be present, as long as this is the only source of Al and Si. When the dissolved Al and Si is then advected into the Atlantic Ocean by the North Atlantic Deep Water (NADW), this signal slowly disappears because of other sources of Al among which dust deposition and possibly sediment resuspension (Middag et al. (2011)) and a source of Si from Antarctic Bottom Water.

Recent years have seen the development of models of the marine biogeochemical cycle of Al. Gehlen et al. (2003) implemented a basic scavenging model, while Han et al. (2008) also included a biological aluminium incorporation module. Gehlen et al. (2003) had the objective of testing the sensitivity of modelled Al fields to dust input and thus to evaluate the possibility for constraining dust deposition via Al_{diss} . To this purpose they embedded an Al cycle in the HAMOCC2 biogeochemical model. The model consists of an equilibrium relation between, on the one hand, Al_{ads} and, on the other hand, Al_{diss} . In chemical equilibrium Al_{ads} is proportional to the biogenic silica (bSiO_2) concentration. In their work, as well as this paper, the term *biogenic silica* or bSiO_2 refers to the detrital fractions which is fuelled by diatoms and other silicifying phytoplankton, which have no stable organic matter coating and sink. When bSiO_2 sinks to the seafloor (together with adsorbed Al), it is buried. The resulting concentration of modelled Al_{diss} was of the same order as the then published observations, but it suggested a significant overestimation of Saharan dust input (Gehlen et al. (2003)) when the dust deposition field of Mahowald et al. (1999) was used. The main goal of Han et al. (2008) was to better constrain the dust deposition field. For this purpose they used the Biogeochemical Elemental Cycling (BEC) model improved by Moore et al. (2008) as a starting point. They used all dissolved Al datasets used by Gehlen et al. (2003) and added more datasets. Except for scavenging Han et al. (2008) added a biological Al uptake module where the Al:Si uptake ratio is a function of the ambient Al and Si concentrations (Han et al. (2008)). The residence time of Al for both modelling studies varies strongly between different locations (from less than one year to almost 200 years), but generally the values are significantly lower than early estimates of 100–200 years (Orians and Bruland (1985)).

Overall, there are a number of questions regarding the oceanic Al cycle that re-

main to be fully addressed. These touch on issues of ocean circulation, the specific sources and sinks of Al in different parts of the world ocean and what processes are needed to accurately simulate the oceanic distribution of Al. Firstly, there is the question of the meridional (north to south) distribution of Al through the Atlantic Ocean. In the North Atlantic Ocean and northern seas, water sinks and forms NADW, which is then transported southward (e.g., Lozier (2010); Gary et al. (2011)). In the deep Atlantic Ocean dissolved silicon (Si) concentration increases from north to south (Ragueneau et al. (2000); Sarmiento and Gruber (2006)), while the concentration of Al_{diss} stays relatively constant until about 20°S and then decreases (R. Middag, in preparation; see also Section 3). Thus it has a generally opposite behaviour compared to Si. Since there are strong suggestions that the processes controlling the distribution of Si and Al are linked, the question is raised how this negative correlation is possible. Secondly, there is the question about the observed profiles of Al_{diss} at different locations in the ocean. Generally, profiles of Al_{diss} have a reversible-scavenging profile (increasing with depth) and often with a minimum near 1 km depth and a maximum at the surface because of dust deposition. However, observations in the Mediterranean Sea (Hydes et al. (1988); Chou and Wollast (1997)) and IPY-Geotraces-NL observations in the eastern Arctic (Middag et al. (2009)) show that there is a strong positive relation between aluminium and silicon. This supports the hypothesis of biological incorporation of aluminium into the cell wall of diatoms (MacKenzie et al. (1978); Stoffyn (1979)).

These issues can be analysed further by the use of numerical models. Since there is a strong spatial variation in aluminium concentration (and its relation to silicon), an ocean general circulation model should be used to simulate the distribution of Al_{diss} . Potentially crucial parameters and sources can be modified in the model to test its sensitivity to these changes. In this way a better understanding of the aluminium cycle can be reached.

In this paper we try to model the observed distribution of Al and examine the processes driving it. Based on new observations and previous work on aluminium modelling (Gehlen et al. (2003), Han et al. (2008)) we formulate a model of aluminium based on dust deposition and scavenging by biogenic silica. This model and the configuration of the simulations will be set out in the following section. Then the observations which we used to check and improve the model will be discussed. The results of the several experiments follow in Section 3, as well as a comparison with the observations. The discussion in Section 4 comprises of a comparison between our model results and Gehlen et al. (2003), a timescale analysis and based on that a discussion of our simulations. We have not compared our results with Han et al. (2008), since we have not performed simulations with biological aluminium incorporation. The last section (5) sums up the conclusions, gives an outline for further development of the model and suggests what further study is needed.

2 Methods

2.1 Model description

To model the three-dimensional distribution of Al_{diss} , the Ocean General Circulation Model (OGCM) called Nucleus for European Modelling of the Ocean (NEMO) is used (Madec (2008)). For this study we use PISCES, one of the biogeochemical components available in NEMO (Aumont and Bopp (2006); Ethé et al. (2006)), which has been employed for many other studies concerning trace metals, as well as large scale ocean biogeochemistry (Aumont and Bopp (2006); Dutay et al. (2009); Arsouze et al. (2009); Dutay et al. (2009); Tagliabue et al. (2010)). PISCES is run off-line forced by a climatological year of physical fields including subgrid turbulence.




Figure 1: Velocity at surface, yearly average, on the ORCA2 model grid. The colour represents the speed in m/s, while the vectors represent the direction of the flow. Vectors are plotted at every fourth gridpoint.

158 The forcing frequency was set to five days. These physical fields were calculated
 159 by the physical component of NEMO named OPA (Madec et al. (1998)), which
 160 was forced by satellite derived wind stress data. Details on the forcings and model
 161 configuration can be found in Arsouze et al. (2009) and Dutay et al. (2009).

162 All input and output fields are defined on the ORCA2 grid, an irregular grid
 163 with a nominal resolution of $2^\circ \times 2^\circ$, with the meridional resolution increased near
 164 the equator and two ‘north poles’ in Canada and Russia to eliminate the coordinate
 165 singularity in the Arctic Ocean. In Fig. 1 the annually averaged surface velocity
 166 field of the OPA output is plotted on the model grid. It shows the main features of
 167 the ocean circulation such as the equatorial current systems, the western boundary
 168 currents (Gulf Stream and Kuroshio) as well as the Antarctic Circumpolar Current.

169 2.1.1 Biogeochemical model

170 We implemented the aluminium cycle (see Section 2.1.2) to PISCES, a biogeo-
 171 chemical model simulating the cycle of carbon and the major nutrients (nitrate,
 172 phosphate, ammonium, silicic acid, and iron), along with two phytoplankton types
 173 (diatoms and nanophytoplankton), two zooplankton grazers (micro- and meso-
 174 zooplankton), two classes of particulate organic carbon (small and large) of dif-
 175 ferential labilities and sinking speeds, as well as calcite and biogenic silica. The
 176 standard version of PISCES accounts for 24 tracers. For a more detailed descrip-
 177 tion of PISCES see the auxiliary material of Aumont and Bopp (2006).

178 The aluminium model interacts with the silicon cycle. The model PISCES dis-
 179 tinguishes three silicon pools: the silicon content of living diatoms (diatom Si); the
 180 silicon content of dead, sinking diatoms (biogenic Si) and dissolved silicic acid. In
 181 the model silicic acid and other nutrients are supplied to the ocean by means of
 182 atmospheric dust deposition, rivers and sediment mobilisation (Aumont and Bopp
 183 (2006); Gehlen et al. (2007)).

184 2.1.2 Aluminium model

185 For the modelling of Al, we introduced two new tracers to PISCES: dissolved alu-
 186 minium (Al_{diss}) and adsorbed aluminium (Al_{ads}). Our Al model follows the ap-

proach of Gehlen et al. (2003), i.e., the internal processes (described below) of adsorption and desorption are identical to the model of Gehlen et al. (2003), although the associated biogeochemical model is more complex.

For the standard model configuration, the sole source of Al to the ocean is by dissolution of dust particles in the ocean surface. In one of the sensitivity experiments, a small fraction is also dissolved in the water column (see Section 2.1.4). We assume an Al:Fe dust fraction of 8.1:3.5, which is based on the mass percentages of Al and Fe known to be present in the Earth’s crust (Wedepohl (1995)). The solubility of Al from dust is not well constrained, with reported values ranging from 0.5–86%, but is probably in the range of 1–15% (Orians and Bruland (1986); Measures and Vink (2000); Jickells et al. (2005)). Furthermore, it is likely that the source of dust has an impact on the solubility of the Al fraction (Baker et al. (2006); Measures et al. (2010) and references herein). We take the dissolution of dust as a constant with respect to the origin and location of deposition. The dissolution occurs only in the upper model layer, and is described by the following equation:

$$\left. \frac{\partial}{\partial t} [\text{Al}_{\text{diss}}] \right|_{\text{deposition}} = \alpha \mathcal{D}_{\text{Al}} / H_1, \quad (1)$$

where $\mathcal{D}_{\text{Al}}$ is the Al flux into the ocean, α is the fraction of Al that is dissolved, and H_1 is the thickness of first model layer. Square brackets denote the concentration of the tracer. Since in the mixed layer has rapid vertical mixing (ν_E is very large in the ocean model; the typical mixing time is D_{ML}/ν_E), adding Al_{diss} in the first layer is effectively the same as adding it spread out over the mixed layer. The Al that does not dissolve from dust, either at the surface or in the water column, is assumed not to play any role in the biogeochemical cycle of Al and on our timescales of interest, and can be thought of as being buried in marine sediments.

Dissolved Al is assumed to be of the form of aluminium that may adsorb onto biogenic silica particles and, aside from external inputs, the Al_{diss} concentration is essentially governed by adsorption and desorption. The Al_{diss} and Al_{ads} concentrations follow the following reversible first-order adsorption equation (Gehlen et al. (2003)):

$$\left. \frac{\partial}{\partial t} [\text{Al}_{\text{ads}}] \right|_{\text{ad/desorption}} = \kappa ([\text{Al}_{\text{ads}}^{\text{eq}}] - [\text{Al}_{\text{ads}}]), \quad (2)$$

where

$$[\text{Al}_{\text{ads}}^{\text{eq}}] = k_d \cdot [\text{Al}_{\text{diss}}] \cdot [\text{bSiO}_2], \quad (3)$$

in which bSiO_2 stands for biogenic silica. $[\text{Al}_{\text{ads}}^{\text{eq}}]$ is the chemical equilibrium concentration of Al_{ads} . The parameter k_d is the partition coefficient and κ is the first order rate constant for equilibration of Al_{ads} to $\text{Al}_{\text{ads}}^{\text{eq}}$. Since total Al is conserved when only internal processes are concerned, the time derivative of Al_{diss} is equal to the negative of the time derivative of Al_{ads} .

Eq. 3 describes the chemical equilibrium between Al_{diss} and Al_{ads} , and Eq. 2 illustrates that some time is needed before equilibrium is reached (it is not modelled as an instantaneous process). As noted by Gehlen et al. (2003), this is a purely empirical model and a more mechanistic description of the adsorption/desorption of Al_{diss} onto bSiO_2 is difficult, since in reality, the pool of observed “dissolved aluminium” is operationally defined, i.e., it exists as different kinds of ions as well as colloids that can go through a filter of, e.g., $0.2 \mu\text{m}$. Also the adsorbent, bSiO_2 , exists in various forms and it is, for instance, difficult to even define the charge density, which is important for its potential to adsorb Al_{diss} . Usually the surface is covered by natural organic matter (Lead and Wilkinson (2007); Filella (2007)). Furthermore, the main scavengers for aluminium in the open ocean also include Particulate Organic Matter (POM), calcium carbonate (CaCO_3) and even particulates

that are not part of PISCES (such as lithogenic particles that do not participate actively in biogeochemical cycles). Hence it is difficult to conceive a mechanistic model of ad- and desorption of “Al_{diss}” onto the poorly-defined pool of particulates.

In our model we use only bSiO₂ as scavenger of Al_{diss}. Biogenic silica is likely to be the most important because (1) it sinks quickly, so that Al has little chance to significantly desorb from bSiO₂ (ballast effect), and (2) positively charged aluminium ions are easily adsorbed onto the negatively charged surface of bSiO₂ particles (Loucaides et al. (2010)). While particulates are also presumed to be negatively charged, observational studies confirm that biogenic silica is a strong scavenger of aluminium (Stoffyn and Mackenzie (1982); Moran and Moore (1989)).

Sinking of bSiO₂ and Al_{ads} is described as follows. In the model, bSiO₂ sinks with a speed of $w_{ML} = 30$ m/day in the mixed layer. Below the mixed layer, it increases linearly with depth, reaching a value of $w_{4km} = 200$ m/day at 4 km below the mixed layer (Aumont and Bopp (2006)). This means that Al_{ads} also sinks with this speed, while it can desorb following equations 2 and 3. For every layer in the model, we have the following equation:

$$\left. \frac{\partial}{\partial t} [Al_{ads}] \right|_{\text{sinking}} = - \frac{\partial}{\partial z} (w_s \cdot [Al_{ads}]) , \quad (4)$$

where w_s is the sinking speed, given in the mixed layer by a constant w_{ML} and below the mixed layer increasing with depth according to:

$$w_s = w_{ML} + (w_{4km} - w_{ML}) \cdot \frac{z - D_{ML}}{4000m - D_{ML}} , \quad (5)$$

where z is the model depth. Once sunk to the ocean floor, we assume that the aluminium is buried permanently. This is described by the following change of Al_{ads} in the bottom layer: $\left. \frac{\partial}{\partial t} [Al_{ads}] \right|_{\text{sedimentation}} = -[Al_{ads}] \cdot w_s / H_{sed}$, where H_{sed} is the thickness of the bottom model layer.

Adding advection and mixing, the full equations for [Al_{ads}] and [Al_{diss}] away from the boundaries are as follows:

$$\frac{d}{dt} [Al_{ads}] = \kappa ([Al_{ads}^{eq}] - [Al_{ads}]) - \frac{\partial}{\partial z} (w_s \cdot [Al_{ads}]) + (\mathcal{A} \nabla_H^2 + \nu_E \frac{\partial^2}{\partial z^2}) [Al_{ads}] , \quad (6)$$

$$\frac{d}{dt} [Al_{diss}] = - \kappa ([Al_{ads}^{eq}] - [Al_{ads}]) + (\mathcal{A} \nabla_H^2 + \nu_E \frac{\partial^2}{\partial z^2}) [Al_{diss}] , \quad (7)$$

where the general advection ($\mathbf{v} \cdot \nabla$) of the tracers is implicit in the full time derivatives. $\mathcal{A}$ and ν_E are the horizontal and vertical eddy diffusivity coefficients, respectively.

The code of the NEMO model, as well as PISCES and the aluminium model, is free software: software that can be used, studied, and modified without restriction (except that published modifications must fall under the same license). It is available at <http://www.nemo-ocean.eu/>. We used version 3.1 of the NEMO model, specifically svn revision 1183. This can be accessed through <http://forge.ipsl.jussieu.fr/igcmg/svn/modips1/>. The aluminium specific code is available under the same conditions as an electronic supplement to this article: `trcal_sed.F90` describes aluminium deposition into the ocean and into the sediment, and `trcal_rem.F90` describes the internal processes of ad- and desorption.

2.1.3 Input fields

The dust deposition field was taken from the output of the INCA model, an atmospheric dust model (Whitehead et al. (1998)). INCA is the aerosol module of the LMDzT atmospheric model (Schulz et al. (2009)). The resulting climatology is




Figure 2: Aluminium sources used to force the model, based on the Fe input in PISCES, using an Al:Fe crustal ratio of 8.1:3.5.

Basin:	Absolute (Gg/yr)	Relative (%)
Atlantic Ocean	4.49	47.6
Indian Ocean	2.08	22.0
Pacific Ocean	1.90	20.1
Southern Ocean	0.013	0.14
Arctic Ocean	0.050	0.57
Mediterranean Sea	0.621	6.58
other basins	0.279	2.96
<i>total</i>	9.436	100.00

Table 1: Amount of yearly dust deposition in each basin. The Southern Ocean is defined as the ocean south of 59°S , without overlap with the Pacific, Atlantic and Indian Oceans.

used and described by Aumont et al. (2008) and evaluated in Textor et al. (2006). The surface deposition is shown in Fig. 2(a). Most dust is deposited in the Atlantic Ocean, just west of the Sahara and just west of the Kalahari Desert. Another important dust deposition sites is found in the northern Indian Ocean. Dust deposition east of Asia in the Pacific Ocean is much smaller, and in the South Pacific Ocean, the Southern Ocean and the Arctic Ocean there is almost no deposition of dust. The amount of dust per basin is listed in Table 1.

According to Textor et al. (2006) the model that is used to create this dust deposition field is validated against high-quality observational datasets, but there are a number of uncertainties, because there are not enough observations available to sufficiently constrain any dust model globally. The spatial resolution is nominally 2° , and the temporal resolution is one month (twelve time steps per year). In reality, dust deposition is highly sporadic (Aumont et al. (2008); Rijkens et al. (2008)). We expect that dust events are highly local and there is sufficient horizontal mixing to dilute the Al_{diss} distribution, such that there is no significant concentration change. Furthermore, our aim is to look at averages of $[\text{Al}_{\text{diss}}]$ over at least several months, and therefore it is acceptable to use monthly averaged dust deposition fields. The deposition field (Fig. 2(a)) is very similar to the one used by Jickells et al. (2005), except that in our field there is almost no dust deposition near Argentina and Chili.

In Fig. 2(b) the sediment source is plotted, which is used for one of the sensitivity experiments. It is a function developed for Fe fluxes and depends on the sea floor topography (which accounts for the degree of oxygenation of the sediments), as described by Aumont and Bopp (2006). The Al:Fe stoichiometry used is 8.1:3.5, the same as for the dust deposition field.

Experiment:	surf.diss. (%)	subsurf.diss.	k_d (10^6 l/kg)	κ (yr^{-1})	sediments
Gehlen et al. (2003)	3	no	1	10000	no
Han et al. (2008)	5	no	-	-	no
Reference experiment	5	no	4	10000	no
Doubled dissolution	10	no	4	10000	no
With subsurface diss.	5	yes	4	10000	no
With ocean margins	5	no	4	10000	yes
Halved part. coeff. k_d	5	no	2	10000	no
Slow equilibration κ	5	no	4	100	no

Table 2: Overview of reference and sensitivity experiments. Deviations from the reference experiment are in bold.

2.1.4 Simulations

Our goal in this study is to better understand how the assumptions made in our model impact on the distribution of Al_{diss} . Specifically, we aim to examine the importance of different Al cycle processes and external Al_{diss} sources. To that end, we perform five sensitivity experiments (see Table 2) with parameters different from the reference experiment.

For the reference experiment we use a five percent dissolution of the Al fraction, which is within the known range and close to the values used by Gehlen et al. (2003) (3%) and Han et al. (2008) (5%). For the partition coefficient k_d we use $4 \cdot 10^6$ l/kg. This quantity signifies the amount of Al that can adsorb onto biogenic silica particulates, which means that if k_d is to be increased, Al_{diss} is exported in larger quantities by scavenging. Our value is four times higher than that of Gehlen et al. (2003), but much smaller than the one used by K. Orians and J. McAlister ($5.6 \cdot 10^7$ l/kg, used for shelves as well as open ocean regions, McAlister and Orians (2011)).¹ The first order rate constant κ , which signifies how quickly the equilibrium between Al_{diss} and Al_{ads} is established, is not constrained by the literature. All these parameters for the reference experiment are chosen in such a way that the model simulates the Al_{diss} distribution from the observations of the selected cruises (see Section 2.2) reasonably well, while all the parameters remained within literature ranges.

After 600 yr the reference experiment is forked into several sensitivity simulations. These sensitivity experiments are run, along with the reference experiment, for another 1300 yr or more to a steady state. The total model time for the reference simulation is 2200 yr. An overview of all key parameters for the simulations is given in Table 2. For all experiments we use the OPA physical fields and atmospheric dust deposition as described in Section 2.1.3. In all simulations, except for the third which tests the sensitivity of water column dissolution, Al is dissolved only in the surface ocean. In the third experiment Al is dissolved in the water column as well with the fraction $10^{-4} \cdot e^{-z/z_0}$ of the dust deposition field, where z is the depth in km and $z_0 = 1$ km. The extra term added to Eq. 7 is $\tilde{\alpha} \cdot e^{-z/z_0} \mathcal{D}_{\text{Al}}/H_k$, where H_k is the thickness of layer k .

The sensitivity simulations can be divided into two types of sensitivity experiments: the first three test the sensitivity to Al sources (amount of dust dissolution into the surface ocean and into the ocean water column, and the inclusion of ocean sediment dissolution) and the last two test the sensitivity to the internal Al cycling (lower partition coefficient k_d and lower first order rate constant κ).

¹We did not use the k_d value from Gehlen et al. (2003), because initial experiments showed that this would leave too much Al_{diss} in the ocean surface, especially at high latitudes.

Cruise	Research vessel	Year	Location	Source	Nr. of data
ARK XXII/2	Polarstern	2007	Arctic	Middag et al. (2009)	1080
ANT XXIV/3	Polarstern	2008	Southern Ocean	Middag et al. (2011, 2012)	919
64 PE 267	Pelagia	2007	North-east Atlantic	Middag et al.	137
64 PE 319	Pelagia	2010	North-west Atlantic	Middag et al.	383
64 PE 321	Pelagia	2010	Centre-west Atlantic	Middag et al.	504
JC057	James Clark Ross	2011	South-west Atlantic	Middag et al.	432
Subtotal used primarily for detailed comparison and optimisation of the model					3455
Compilation of selected other observations for global ocean comparison:					
IOC96	Knorr	1996	Central-south Atlantic	Vink and Measures (2001)	1049
M 60	Meteor	1982	North-east Atlantic	Kremling (1985)	91
IRONAGES III	Pelagia	2002	North-east Atlantic	Kramer et al. (2004)	181
EUCFe	Kilo Moana	2006	Equatorial Pacific	Slemons et al. (2010)	195
MC-80	Thompson	1980	Pacific	Orians and Bruland (1986)	92
VERTEX-4	Wecoma	1983	North Pacific	Orians and Bruland (1986)	54
VERTEX-5	Thompson	1984	North Pacific	Orians and Bruland (1986)	59
KH-98-3	Hakuho-Marui	1996	East Indian	Obata et al. (2007)	152
Subtotal compilation of selected other observations for global comparison					1873
Grand total of all dissolved Al values used in this study					5328

Table 3: Observational data used for comparison with model. For positions see Fig. 3.

2.2 Observations

The recent IPY-Geotraces observations in the Arctic (Middag et al. (2009)), North-east Atlantic (Middag (2010), Chapter 5), West Atlantic (R. Middag, in preparation) Oceans and the Atlantic section of the Antarctic Ocean (Middag et al. (2011)) are used for a detailed comparison and optimisation of the model parameters. See the upper part of Table 3 for these datasets and Fig. 3 for the station positions. These datasets comprise overall 3455 individual data values for dissolved Al. All values have been verified versus international reference samples and their consensus values of the SAFe and Geotraces programmes, see supplementary material S-1. Moreover, during cruise 64 PE 321 (2010) excellent agreement was obtained for a complete vertical profile of dissolved Al at the BATS station between a previous occupation and dissolved Al data in 2008 by the US Geotraces group, see supplementary material S-2. This large amount of observations give the possibility to compare the model with deep ocean observations of $[Al_{diss}]$, allowing us to validate the model in the deep ocean and to study the global Al cycle in more detail.

For a worldwide global ocean comparison one has to rely on data that was collected in the era before the reference samples of SAFe and Geotraces were available. Inevitably the definition of criteria for selecting such previously published datasets is less rigorous, see Electronic supplement S-3 for the criteria used for each of the selected datasets. The selected datasets are listed in Table 3, lower part, and positions are shown in Fig. 3.

3 Results

3.1 Reference simulation

Our reference experiment is performed for a spin-up period of 2200 yr. The resulting total ocean Al budget in the reference simulation is around 7 Tmol (1 Tmol = 10^{12} mol). Already after 600 yr the total Al distribution is more or less in a steady state




Figure 3: Stations of the cruises: M 60 [yellow]; MC-80 and VERTEX [red]; IOC96 [green]; IRONAGES III [blue]; KH-98-3 [pink]; EUCFe [black]; Geotraces-NL [purple]. See Table 3 for an overview with references and the number of observations.

as shown by Fig. 4, where the total integrated aluminium (dissolved and adsorbed) is plotted against time. Therefore, from 600 yr our sensitivity experiments (see Section 3.2) are forked.²

Fig. 5 shows the Al_{diss} surface concentration of the reference experiment. As in all subsequent plots, yearly averages are shown. There is a seasonal cycle in $[\text{Al}_{\text{diss}}]$ in our model, but it is small relative to the spatial trends. The observed Al_{diss} concentrations from our Al data compilation (see Section 2.2) are plotted as circles over the model Al_{diss} concentration in this and later figures. A comparison of the model with the observations will be done in Section 3.1.1. Consistent with the dust deposition field (Fig. 2(a)), the largest modelled Al_{diss} surface concentrations are in the central Atlantic Ocean, with values between 30 and 40 nM ($1 \text{ nM} = 10^{-9} \text{ mol/l}$) near 20°N , decreasing northward to values in the order of several nM north of 60°N . The concentration of Al_{diss} between 40°N and 60°N is larger than expected based on the dust deposition field (Fig. 2(a)). This is because the Gulf Stream and the North Atlantic Current transport Al_{diss} northward before most of it is scavenged. As we will see in Section 4.2.1 and 4.5, most Al_{diss} is scavenged before it passes Iceland. The result is that $[\text{Al}_{\text{diss}}]$ in the surface waters of the Arctic Ocean is in the order of 1 nM. On the other side of the globe, south of 40°S , the Al concentration is even lower, less than 1 nM. This is to be expected on the basis of $[\text{bSiO}_2]$ (Fig. 6) and the dust deposition flux (Fig. 2(a)). In the Pacific Ocean the concentration is relatively low as well, except near $20\text{--}35^\circ\text{N}$ where a wide band of high $[\text{Al}_{\text{diss}}]$ can be found. This maximum arises from the large dust deposition along the Asian continental margin between 20°N and 60°N . North of 40°N , the concentration of bSiO_2 is high, as can be seen in Fig. 6. Therefore, in that region the dissolved Al is quickly depleted by the high bSiO_2 concentration in the surface, while south of 40°N the North Pacific Current transports it eastward into the central North Pacific Ocean.

In Fig. 7 the modelled Al_{diss} concentration of the reference experiment is shown at selected depths (100 m, 500 m, 1000 m and 3000 m) and in Fig. 8 the Geotraces cruise section in the West Atlantic Ocean. This section was calculated from the

²The relevant tracers of the raw model output can be found at http://data.zkonet.nl/index.php?page=Project_view&id=2916&tab=Datasets.




Figure 4: Total Al budget ($[Al_{ads}] + [Al_{diss}]$) (Tmol) in the world ocean, plotted against the model years




Figure 5: Dissolved aluminium concentration (nM) from the reference experiment at the surface ocean, with respective observations plotted on top of it. Note that the scale is expanded for low concentrations.




Figure 6: Biogenic Si concentration (nM) in the PISCES model, at the surface, yearly average.




Figure 7: Dissolved aluminium concentration (nM) from the reference experiment at several depths, with respective observations plotted on top of it.




Figure 8: Dissolved aluminium (nM) from reference experiment along the West Atlantic Geotraces cruise track. Observations are plotted on top.

three-dimensional model data by converting the ORCA2 gridded model data to a rectilinear mapping and after this interpolating the rectilinear data onto the cruise track coordinates. In the Atlantic Ocean there is a maximum of $[Al_{diss}]$ from about 30°S at 2 km depth, getting higher northward (Fig. 8). According to the model, deep Al_{diss} concentrations are low in the Southern Ocean, the Arctic Ocean and the Pacific Ocean (Fig. 7).

3.1.1 Comparison with observations

In Figs. 5, 7 and 8 observed Al_{diss} concentrations (see Section 2.2) are plotted as circles over the modelled results. The precision of the measurements performed during the Geotraces cruises at a 1 nM level and lower (polar oceans) was about 4% and at a 20 nM level (West Atlantic Ocean) about 2% (Middag (2010), p. 25). These observations have a standard deviation that is smaller than can be distinguished based on the colour bar. Older observations generally have lesser precision, but of sufficient quality (see Section 2.2). Therefore the model results can in principle be compared with the observations without regard of measurement errors.

Fig. 5 shows that the model and the observations reveal similar patterns in the surface ocean: high concentrations in the central Atlantic Ocean and low concentrations in the polar oceans and the Pacific Ocean. However, details in the modelled Al_{diss} distribution do not agree with the observations. For instance, the large concentration in the Pacific Ocean between 20 and 35°N is not visible in the observations of Oriens and Bruland (1986). Also between 45 and 65°N in the West Atlantic Ocean the concentration of Al_{diss} in the model is overestimated compared with the Geotraces observations near Greenland (R. Middag et al., in preparation). This suggests that Al is not sufficiently scavenged in this area. This is further discussed in Section 4.2. The model underestimates $[Al_{diss}]$ near Brazil. This can be related to errors in ocean currents (Fig. 1), as the transport of low $[Al_{diss}]$ water might be too high in the Brazil current. In reality this current is narrower than the 2° resolution of the model. Another possibility is that the dust deposition field is not realistic in this area. Furthermore, the observations generally have more spatial variability than the modelled $[Al_{diss}]$. Many of these differences depend on several model boundary conditions (dust deposition distribution, velocity field, particle sinking speed) as well as temporal variability, because the model results plotted are yearly averages, while observations are done throughout the year in




Figure 9: The quantity in the coloured area is the model $[Al_{diss}]$ (nM) at the Geotraces West Atlantic cruise section. The contours represent the model Atlantic Overturning Stream Function (Sv).

precise locations.

Fig. 8 shows that similar patterns are present in the West Atlantic Ocean in the model and the observations. There is a clear pattern of North Atlantic Deep Water (NADW), which is mainly visible between 40°S and 20°N in both the model and the observations. In Fig. 9 the Atlantic Overturning Stream Function (OSF) is plotted over the model results, showing that the patterns of the OSF and $[Al_{diss}]$ coincide. In the Southern Ocean, low concentrations of Al_{diss} in the upper km are clearly penetrating from the Southern Ocean northward to at least 20°S around 500 m depth. This is a mixture of Antarctic Intermediate Water (AAIW) and Subantarctic Mode Water (SAAMW). From the Southern Ocean near the bottom, there is also Antarctic Bottom Water (AABW) flowing northward, visible in the observations and the model.

The similarity between the model and the observations lessens in the deeper North Atlantic Ocean, where according to the observations $[Al_{diss}]$ increases with depth (for depths below 800 m), while in the model there is a decrease with depth below 1.5 km. Besides this general pattern of increase of $[Al_{diss}]$ with depth in the observations, a very high concentration of Al_{diss} is present between 45 and 50°N near the sediment, which enhances the dissimilarity between the model and the observations. This problem will be discussed further in Section 4.5.

Furthermore, in the southern hemisphere in the model the maximum of $[Al_{diss}]$ is at about 2 km depth, indicating a southward flow around 2 km depth. In the observations this maximum is at 3 km depth, indicating a southward flow at 3 km depth. Indeed, according to the hydrographical analysis during the West Atlantic Geotraces cruise (Van Aken et al.) the flow is at 3 km depth. As can be seen in Fig. 9 the southward flow is too high in the water column compared to the observations.

Another feature which is not captured by the model is the aluminium profile in the eastern Arctic Ocean, which is shown in Fig. 10. It shows that the model $[Al_{diss}]$ increases with depth from a minimum near 150 m to a maximum near the bottom by a factor of 3 (and is in the order of 2 nM, see Fig. 10). The observations show an increase as well (as also presented in Fig. 7), but here it is far more pronounced with a small minimum at the surface (less than 1 nM) to a large maximum near the abyss (around 18 nM) (Middag et al. (2009)). Thus in the Arctic Ocean the vertical change in $[Al_{diss}]$ in the model is by far not as pronounced as in the observations. The absence of sediment sources of Al might play a role in this. The results of a simulation with sediment sources will be discussed in Section 3.2.




Figure 10: $[Al_{diss}]$ (nM) profile in the eastern Arctic of reference experiment (blue dashed line) and the observations of IPY-Geotraces-NL observations (black solid line) at 213.6°E, 87.03°N

3.2 Sensitivity simulations

Section 2.1.4 described how our sensitivity simulations are set up. The reasons why we performed these specific experiments are closely related to the problems we encountered while interpreting the results from the reference experiment (Section 3.1). After a brief summary of two of these problems and associated sensitivity simulations, we will present the results of these simulations.³

How can it be that in the Atlantic Ocean from 800 m depth to the bottom of the ocean there is a decrease in the Al_{diss} concentration in the model, while according to the observations there is an increase (see Fig. 8)? The only transformation mechanism in our model is reversible scavenging, which should give a profile where $[Al_{diss}]$ increases with depth, and the only source is dust. However, the model biogenic silica concentration, its sinking speed, the physics and the first order rate constant κ are all relevant for the vertical distribution of Al_{diss} as well. Since the biogenic silica concentration and the physical fields can both be assumed to be reasonably realistic (see Section 2.1), no sensitivity simulations concerning these fields are described in this paper. We do perform one simulation with a different first order rate constant (Section 3.2.5) to investigate where in the ocean this parameter influences the Al_{diss} distribution. The value of κ is significantly decreased in this simulation (this parameter is very poorly constrained by observations).

Concerning the strongly depth-increasing profile of Al_{diss} as observed in the eastern Arctic Ocean (Fig. 10), we perform a simulation where sediment ocean margins are added as a source of Al_{diss} , following the approach of Aumont and Bopp (2006) and assuming an Al:Fe fraction of 8.1:3.5 in the sediments (thus we assume sediment Al_{diss} supply is coupled to Fe). The result of this simulation is

³The relevant tracers of the raw model output can be found at http://data.zkonet.nl/index.php?page=Project_view&id=2916&tab=Datasets.




Figure 11: The relative difference of the dissolved aluminium concentration between the experiment with an increased dissolution (10%) and the reference experiment (5% aluminium dissolution). The scale is in percentages, of which the largest part from 95% to 100%.

discussed in Section 3.2.3. Furthermore, an experiment is performed where we have halved the partition coefficient from $4 \cdot 10^6$ l/kg to $2 \cdot 10^6$ l/kg, which is a way to evaluate the importance of relative partitioning between the dissolved and adsorbed fraction. This experiment will be discussed in Section 3.2.4.

3.2.1 Increased dust dissolution fraction

To examine the impact of greater dust dissolution on the cycling and distribution of Al_{diss} , we increase the surface dissolution percentage of Al from 5 to 10% (‘doubled dissolution’). In Fig. 11 the resulting change in $[\text{Al}_{\text{diss}}]$ in the surface ocean as well as in the West Atlantic Ocean (along the Geotraces cruise track) is presented. The $[\text{Al}_{\text{diss}}]$ is increased with a factor of two in most locations in the ocean.

The doubling can be explained when the model equations (Eqs. 2 and 3) are considered. Consider a one-box model in which equilibration is instantaneous ($\kappa = \infty$). If we increase dust dissolution everywhere with a factor of two and wait until steady state, we also get a doubled particulate Al burial. This means that $[\text{Al}_{\text{ads}}]$ must be doubled in the ocean since the sinking/burial speed does not change. Since in steady state $[\text{Al}_{\text{ads}}]$ is proportional to $[\text{Al}_{\text{diss}}]$ (Eq. 3), $[\text{Al}_{\text{diss}}]$ must be doubled as well. Hence a doubled $[\text{Al}_{\text{diss}}]$ is expected. In other words, the total Al budget is linear with the input. (See for instance Broecker and Peng (1982) for background.) This means that the amount of dissolution in the surface ocean has an effect everywhere in the ocean, i.e. the effect is global and not restricted to dust deposition sites.

The Al budget is not doubled everywhere. In Fig. 11(b) we can see that in the coastal upwelling region near Chili the increase is only around 95%. This is because the model is not completely spun up, as can be seen in Fig. 12, so that the increased dissolution of Al from dust has not reached the deep Pacific Ocean yet. Therefore the percentages in Fig. 11 indicate how close the subsequent sensitivity simulations are to steady state in different regions in the ocean.

3.2.2 Water column dust dissolution

For the reference experiment we included dust dissolution only in the surface ocean. to assume that Al from dust only dissolves in the surface ocean. During sinking of dust more Al might be dissolved. To test the effect of water column dissolution we performed a sensitivity simulation with water column dissolution included.

Gehlen et al. (2003) have included Al_{diss} input at the surface only, but they do compare different dust deposition distributions, as do Han et al. (2008). The latter also dissolve dust below the surface layer, but no sensitivity experiments have been




Figure 12: Total Al budget (Tmol) in world ocean during spin-up of 1490 years for increased dissolution sensitivity simulation (solid black line; forked from the reference simulation at year 600). The dashed blue line signifies the Al budget of the reference simulation.




Figure 13: Difference in $[Al_{diss}]$ (nM) between the simulation with water column dissolution with the reference simulation (without water column dissolution)

presented for water column dissolution. None of our sensitivity experiments, as part of an OGCM, have been published before.

Our simulation with water column dissolution resulted in slightly higher values of $[Al_{diss}]$ of up to 0.7 nM more in some places. Fig. 13 compares $[Al_{diss}]$ in the water column dissolution experiment with the reference experiment (Figs. 5, 8).

On first sight, north of the equator, water column dissolution moves the modelled $[Al_{diss}]$ away from the observations. This is clear if we consider the decrease in $[Al_{diss}]$ with depth in the reference simulation, while in the observations $[Al_{diss}]$ increases with depth. The simulation with water column dissolution makes the modelled decrease even more pronounced. Also the high $[Al_{diss}]$ in the upper 500 m is only slightly improved, and not at precisely the right location: in the upper 500 m the model especially underestimates $[Al_{diss}]$ between 20 and 35°N. Therefore, water column dissolution does not improve the model results.




Figure 14: Experiment with sediment input: $[Al_{diss}]$ (a,b) and its difference with the reference run (c,d)

3.2.3 Ocean sediments source

It is possible that dissolution of dust is not the only way in which aluminium enters the ocean. We excluded river input and hydrothermal vents as significant sources of Al (see Section 1). But one other source that might be significant, and might also help our model to produce more realistic $[Al_{diss}]$ (e.g. in the Arctic Ocean), is redissolution of Al from sediment resuspension. Therefore, in one of the simulations, we have included a rough approximation of Al margin sources, as described in Section 2.1.3. The resulting $[Al_{diss}]$ and its difference with the reference experiment is plotted in Fig. 14. We can clearly see the influence of shelf sediment sources on the aluminium distribution. Especially in the Arctic Ocean, the Indonesian Archipelago and near the east coast of North America, dissolved aluminium is increased substantially when sediment input is added to the model. This is to be expected, since the largest sediment source is on the shelf areas (see Fig. 2(b) for the added Al source).

Let's now look at the West Atlantic cruise track. Fig. 14(c) represents the difference between the sediment source experiment and the reference simulations at the ocean surface. There is a clear increase at and near the surface between 20 and 40°N in the West Atlantic Ocean. The cruise section difference plot in Fig. 14(d) shows a large increase in the NADW. So it seems that high $[Al_{diss}]$ can be explained by not only dust deposition but also sediment sources. This is consistent with the explanation of Middag et al. (in preparation), even though their arguments mostly concern deep sediment resuspension and to a lesser extend margin sediments. The sediment source in our model is mostly from near-shore sediment resuspension (Fig. 2(b)), in line with the findings of Mackin and Aller (1986). Of course, a deep sediment source might also explain the high $[Al_{diss}]$ near the sediment at 45–50°N, and it may therefore be important to include such a source in a future model study, since our current results suggest that margin sediments do not effectively reproduce this maximum.




Figure 15: $[\text{Al}_{\text{diss}}]$ (nM) profile from a station in the eastern Arctic (ARKXXII/2): observations (black solid line), the reference simulation (blue dashed line), and sediment sensitivity simulation (red dash-dotted line), all at 213.6°E , 87.03°N

556 The concentration in the model gets significantly higher in the Arctic Ocean
 557 which is bad for depths between the surface and a few km, but better for deeper
 558 waters, where a much higher concentration of aluminium was observed. This is
 559 shown in the profile plots of Fig. 15. Sedimentary input has as an effect that $[\text{Al}_{\text{diss}}]$
 560 increases everywhere in the Arctic Ocean and especially with depth, giving a profile
 561 that has the same *form* as the observations, suggesting the potential importance of
 562 margin Al_{diss} sources herein. However, the modelled $[\text{Al}_{\text{diss}}]$ is too large compared
 563 with the observations. Potential reasons for this are described in Section 4.3.

564 3.2.4 Decreased partition coefficient

565 One of the key parameters in our model is the partition coefficient k_d . It regulates
 566 the amount of Al that can be adsorbed by bSiO_2 . To see what effect this parameter
 567 has in our model setup, we performed an experiment in which we decreased k_d by
 568 a factor of two.

569 Fig. 16 shows that a decrease of k_d leads to the most significant relative increase
 570 of $[\text{Al}_{\text{diss}}]$ in the Southern Ocean. The reason for the large increase in $[\text{Al}_{\text{diss}}]$ in the
 571 Southern Ocean, and also a significant amount in the northern seas, is that these
 572 are locations of high diatom production, which results in large $[\text{bSiO}_2]$. As can be
 573 seen from Eq. 3, the equilibrium concentration of Al_{ads} is proportional to k_d and
 574 $[\text{bSiO}_2]$, hence k_d is important in regions with high $[\text{bSiO}_2]$. This will be further
 575 explained by looking at the Al budget and timescales in Section 4.4.

576 3.2.5 Decreased first order rate constant

577 The second most important scavenging parameter is the first order rate constant κ .
 578 This parameter should have a dynamic effect on the distribution of Al_{diss} , since it




Figure 16: Experiment with a decreased k_d with (a,b) $[Al_{diss}]$ (nM), and (c,d) the relative difference with the reference experiment.

describes how quickly Al_{diss} and Al_{ads} equilibrate (see Eq. 3). To test this in our model, we decreased κ from 10^4 to 10^2 yr^{-1} .

As can be seen in Fig. 17, surface $[Al_{diss}]$ increases significantly compared with the reference experiment (compare Fig. 5 with Fig. 17(a)). Fig. 17(d) shows the relative change in $[Al_{diss}]$ in the West Atlantic Geotraces section. From the equator northward below 2 km depth an increase of aluminium is visible. There is also an increase near the deep sediment between 45 to 50°N, but it is too small to explain the elevated concentration in this area found in the West Atlantic Geotraces observations (see bottom right in Fig. 8).

4 Discussion

4.1 Comparison with Gehlen et al. (2003)

Since our model is very similar to the model of Gehlen et al. (2003), we expect a good agreement between our results and theirs. Even though several features in the two model results are the same, there are also noteworthy differences. These can arise because of differences in the ocean physics, the dust deposition field, as well as the biogeochemical model.

For the velocities the output of the NEMO model on the ORCA2 configuration is used in our model. On the other hand, in the model of Gehlen et al. (2003) a climatology from the Hamburg Large-Scale Geostrophic (LSG) OGCM is used (Maier-Reimer et al. (1993)). The basic characteristics of the physics of the two models are the same, but a few features, like the Mixed Layer Depth (MLD) and the Atlantic MOC, are significantly different. Even though LSG is a low resolution model, the depth of the meridional southward return flow of the Atlantic MOC is almost 3 km deep, which is realistic. On the other hand, in the OPA model (with the ORCA2 configuration) this return flow is much more shallow (a bit over 2 km deep), as shown in Fig. 9. A shallow return flow is typical for low resolution models




Figure 17: $[Al_{diss}]$ (nM) in the slow equilibration experiment ($\kappa = 100 \text{ yr}^{-1}$) and the relative difference with the reference experiment ($\kappa = 10^4 \text{ yr}^{-1}$).

(e.g. Dutay et al. (2002)). Furthermore, the AABW in LSG goes to 15°N, while in our model it goes all the way to 40°N. Overall, the physics of the two models are comparable, but we should keep in mind that the overturning in the ORCA2 configuration of OPA is too shallow and AABW goes too far north, compared with the LSG model.

In both the work of Gehlen et al. (2003) and in our work, the highest concentration of Al_{diss} is found in the surface model layer (Fig. 5) in the Atlantic Ocean around 20°N. In both models the Al_{diss} concentration is low in the Southern Ocean and in the Pacific Ocean. However, there are distinct differences of $[Al_{diss}]$ in the surface ocean in the models. Several features visible in the observations are better captured by the model of Gehlen et al. (2003), while other features are simulated better by our model. These difference are likely to be due to the different dust deposition fields used in the model of Gehlen et al. (2003) compared to the one we used.

In the eastern Atlantic Ocean around 20°N near the coast of Africa, Gehlen et al. (2003) simulate a Al_{diss} concentration of over 300 nM, or around 50 nM for a different dust deposition field. The concentrations in our model and the observations, however, are generally around 30 nM.

In and near the Indonesian Archipelago $[Al_{diss}]$ ranges from 1 to 20 nM in both models, while the observations of Obata et al. (2007) and Slemons et al. (2010) are much more homogeneous, about 10 nM. This might be because the ocean currents in both models are not realistic because of a too low resolution of our model for this region. Possibly other effects in the real ocean play a role, like sedimentary input from the Indonesian Archipelago and east of Indonesia (see Fig. 2(b) for the relative importance).

The vertical meridional section in Fig. 8 through the West Atlantic Ocean shows a reasonable correspondence between the two models. The concentration at the surface is high in both models, it roughly decreases with depth in the Atlantic Ocean. In the Southern Ocean aluminium concentrations are low in the whole

column, but especially in the surface (Fig. 8, very left). However, in our model between 10°S and 45°N there is a significant decrease of $[Al_{diss}]$ with depth, which is not visible in the model results of Gehlen et al. (2003) where below 2 km north of the equator $[Al_{diss}]$ is rather homogeneous in both dimensions along the cross-section. The model of Gehlen et al. (2003) shows North Atlantic Deep Water (NADW) in the Al_{diss} distribution, spreading southward at a depth of almost 3 km, as also shown by the observations (Fig. 8, but also Van Aken, in preparation). Our model also captures NADW, but this is at only 2 km depth, as explained before.

4.2 Advection versus scavenging

While Al is entering the ocean through the surface by dust deposition and leaving it at the bottom through burial, it is also transported by the currents. In this section we try to find out whether scavenging or advection is more important in different regions of the ocean, and how this can explain certain features in the model results.

In our model scavenging is reversible, which we base on a priori arguments (Bacon and Anderson (1982); Anderson (2006)) and the general increase of $[Al_{diss}]$ with depth in observations. Therefore it is unexpected that in the Atlantic Ocean a depth decreasing profile of Al_{diss} is simulated (see e.g. Fig. 17(b)). One possible reason for this is that the $bSiO_2$ distribution is not realistic. However, the PISCES model with the Si cycle is used by several modelling studies (Aumont and Bopp (2006); Dutay et al. (2009); Arsouze et al. (2009); Dutay et al. (2009); Tagliabue et al. (2010)) and has the same spatial patterns as known $bSiO_2$ export fields (e.g. Sarmiento and Gruber (2006)).

We should emphasise that we only scavenge Al by biogenic silica. Of course $CaCO_3$ and POC might also scavenge in reality. Looking back to our comparison with the observations of R. Middag et al., we found that near Greenland $[Al_{diss}]$ was overestimated by the model. This would be an example where the addition of other scavengers, especially $CaCO_3$ can help in decreasing the modelled $[Al_{diss}]$ near Greenland, and also in the western South Atlantic Ocean between 30 and 40°S. Refer to e.g. Dittert et al. (2005); Sarmiento and Gruber (2006) for export of $bSiO_2$, $CaCO_3$ and POC; or Lam (2011) for concentrations.

The importance of scavenging is visible from several features in our model data. Firstly, Fig. 7 shows that on a global scale the pattern of $[Al_{diss}]$ has roughly the same form at different depths, which is a result of reversible scavenging, i.e., at the surface where $[bSiO_2]$ is high, Al is mainly adsorbed onto $bSiO_2$, while at depth Al is mostly desorbed from $bSiO_2$. Specifically, $[Al_{diss}]$ is high in the Atlantic and Indian Oceans north of 30°S, and much lower in other regions of the ocean. Secondly, the fact that the concentration in the Pacific Ocean is low at all depths, is because Al_{diss} is removed by scavenging before it reaches the Pacific Ocean. These are just a few of the features in the Al_{diss} distribution that suggest that on a large scale scavenging is more important than advection.

However, the observations and the model show that advective transport by ocean currents is important as well. For instance, Fig. 2(a) shows that large quantities of dust are deposited just west of the Sahara. However, large concentrations of Al_{diss} are not only visible directly below the dust deposition site, but even higher concentrations are found further to the west (Fig. 5). Large amounts of dissolved Al must be advected from the dust deposition site toward Central America. Hereafter it seems to be advected northward until Iceland, where $[Al_{diss}]$ is high despite very little dust deposition. Deeper in the North Atlantic Ocean the original dust signal decreases, and at 2 km depth in the model (or 3 km depth in the observations) Al_{diss} is transported into the southern hemisphere by the NADW. This southward transport can be seen clearly in Fig. 8, but no further than 40°S. South of this latitude the Al_{diss} concentration is low. These examples indicate that also advection

686 plays an important role in the redistribution of Al.

687 4.2.1 Timescales

688 While the analysis of $[Al_{diss}]$ distributions points to the combined importance of
 689 scavenging and advection in setting its overall pattern, it does not allow to quantify
 690 their relative importance. In order to assess the relevance of these different processes
 691 we define associated timescales based on our model equations.

A priori the relative relevance of scavenging can partly be derived from the model equations. If Eq. 3 is substituted into Eq. 2, we get:

$$\begin{aligned} \frac{d}{dt}[Al_{ads}] &= \kappa \cdot ([Al_{ads}^{eq}] - [Al_{ads}]) \\ &= \frac{1}{\tau_{trans}}[Al_{diss}] - \frac{1}{\tau_{eq}}[Al_{ads}] , \end{aligned} \quad (8)$$

692 where $\tau_{trans} = 1/(\kappa k_d [bSiO_2])$ is the typical adsorption timescale. This is the typical
 693 time necessary for Al to transform from Al_{diss} to Al_{ads} (or vice versa). This process
 694 depends on the amount of available $bSiO_2$ and the amount of Al_{diss} which can
 695 adsorb onto $bSiO_2$, giving the actual rate of conversion between Al_{diss} and Al_{ads} .
 696 The addition of Al_{ads} is equal to the removal of Al_{diss} , so τ_{trans} is either the time
 697 necessary to add Al_{ads} , or the time to remove Al_{diss} . The second timescale in Eq. 8,
 698 $\tau_{eq} = 1/\kappa$, is the time it takes for Al_{ads} to equilibrate to Al_{ads}^{eq} . This process provides
 699 a stabilising feedback on the growth of $[Al_{ads}]$ (or decrease of $[Al_{diss}]$).

700 Scavenging is the process of adsorption and sinking, so we need to know how
 701 fast particles sink. We use this together with the adsorption timescale to define a
 702 scavenging timescale. Since we look at the surface ocean, we might define the typical
 703 sinking speed as sinking through the first 10 m, the thickness of the upper model
 704 gridbox. However, after exporting Al_{ads} out of a gridbox by sinking, the particles
 705 sink through the mixed layer (which is generally thicker than 10 m). During sinking,
 706 the particles will partly be mixed back into the upper gridbox. Therefore to speak
 707 of a removal, it is a necessary condition for Al to sink out of (at least) the mixed
 708 layer. In our model, the sinking velocity of $bSiO_2$ and Al_{ads} is defined to be constant
 709 in the mixed layer, namely $w_s = 30$ m/day. Now we are easily able to define the
 710 typical sinking timescale: $\tau_{sink} = D_{ML}/w_s$, where D_{ML} is the mixed layer depth.
 711 The scavenging timescale can be defined as the maximum of the adsorption and the
 712 sinking timescales:

$$\tau_{scav} = \max(\tau_{trans}, \tau_{sink}) . \quad (9)$$

713 This signifies how fast Al is exported from the mixed layer. Except for a few places
 714 where the mixed layer is very deep, like the centre of the Labrador Sea, the time it
 715 takes to sink out of the mixed layer is generally very small compared to the time it
 716 takes the Al to adsorb onto a particle ($\tau_{sink} \ll \tau_{trans}$) and thus almost everywhere
 717 $\tau_{scav} = \tau_{trans}$.

718 A typical advection timescale is defined as follows:

$$\tau_{adv} = L/V . \quad (10)$$

719 Here L is the typical length scale, defined as the horizontal diameter of a gridbox,
 720 and V is the horizontal speed. Since the vertical velocity component is very small
 721 ($W \ll (D_{ML}/L)U$), only the horizontal components need to be considered. The
 722 meaning of this timescale is that within a time τ_{adv} a dissolved Al particle (or any
 723 other non-buoyant tracer) is advected out of a gridbox (if it is not scavenged before).

724 The residence time can be defined as the minimum of the scavenging and the
 725 advection timescales. It is the typical time that a Al_{diss} particle stays within a
 726 volume box of horizontal resolution (about $2^\circ \times 2^\circ$) times D_{ML} . This quantity




Figure 18: Residence time in months of Al in a box of horizontal resolution times D_{ML} , or $\tau_{res} = \min(\tau_{adv}, \tau_{scav})$. It represents the average time an Al_{diss} particles spends in this box, considering only scavenging and horizontal advection.

is presented in Fig. 18. The oligotrophic gyres in the Atlantic, Pacific and Indian Oceans are clearly visible. In the centre of these gyres it takes much longer than one year before Al is exported out of a volume box, either by advection or by scavenging. The modelled Al_{diss} distribution of Fig. 5 shows in the Atlantic Ocean the largest values between 10 and 30°N. This can be partly explained by the large dust input between 5 and 20°N, where Al_{diss} is removed relatively fast by advection but dust input keeps $[Al_{diss}]$ large, and partly by the large residence time between 20 and 30°N, where there is no large dust input but Al_{diss} simply stays there for a long time, since there is no $bSiO_2$ present for scavenging and advection is very small. A similar argument can be given for the high $[Al_{diss}]$ near the North Pacific gyre.

If we want to know which of the timescales are more important, a relative relation between τ_{scav} and τ_{adv} must be defined. The number for relative importance of scavenging versus advection for Al export is the following:

$$\Upsilon = \frac{\tau_{adv}}{\tau_{scav}}. \quad (11)$$

The logarithm of this quantity is plotted in Fig. 19. In regions where $\Upsilon \gg 1$, like north of 40°N in the Atlantic and Pacific Oceans and in the south of the Southern Ocean, scavenging is more important than advection. On large scales the large Υ regions coincide very well with the large $bSiO_2$ regions (Fig. 6). Looking more in detail, we can see that Υ is less homogeneous than $[bSiO_2]$. For instance, in Drake Passage and in the Pacific sector of the Southern Ocean at 150°W the timescale fraction is smaller then one would expect based on the $bSiO_2$ distribution. This is because of the strong current of the Antarctic Circumpolar Current (see Fig. 1).

In regions where $\Upsilon \ll 1$, in the low latitudes, advection is more important than scavenging. One can say that if $\Upsilon \ll 1$, Al export is advection driven, while for $\Upsilon \gg 1$, Al export is scavenging driven. In the high Υ regions a one-dimensional model would be a reasonable approximation, provided that there is sufficient Al input to keep $[Al_{diss}]$ in steady state. One region where this should perfectly work is the Mediterranean – there Υ is very large and the dust flux is high. Several other regions like the northern seas, near Antarctica and some other coastal areas would also be good candidates for a one-dimensional model. Other regions of the ocean cannot be well described by a one-dimensional model. Specifically the Arctic




Figure 19: $\log_{10}(\Upsilon)$, when $\kappa = 10^4 \text{ yr}^{-1}$ and $k_d = 4 \cdot 10^6 \text{ l/kg}$. For the advection time τ_{adv} the definition of Eq. 10 is used, and the scavenging time τ_{scav} is the maximum of τ_{trans} and τ_{sink} .

757 Ocean would not be a good candidate. Even though it is a semi-closed basin,
 758 internal circulation has an effect of the same order as scavenging. These arguments
 759 must be taken into consideration when using one-dimensional models.

760 4.3 Ocean sediments source

761 Including margin sediment input of Al results in overestimated values for $[\text{Al}_{\text{diss}}]$
 762 compared to the observations in the Arctic Ocean. Potential reasons for this will
 763 now be discussed.

764 As described in Section 2.1.3 the Al flux is proportional to the Fe flux which
 765 depends on the degree of oxygenation of the sediments. This is probably not a rea-
 766 sonable assumption, since generally Al does not flux out of undisturbed sediments,
 767 but enters the water column through resuspension of the sediments (e.g. Mackin
 768 (1986); Van Beusekom et al. (1997)). Since there is not enough data on such resus-
 769 pension, we choose this simplistic parametrisation for our sensitivity experiment.
 770 A priori we therefore have no reason to expect an improvement in the model. The
 771 sensitivity experiment can only be used to get an idea of the first order effects of
 772 an ocean sediment source of Al, henceforth referred to as “sediment input”.

773 Since the sediment input in the Arctic Ocean is treated in the same way as the
 774 rest of the ocean, it is surprising that $[\text{Al}_{\text{diss}}]$ gets much too high (Fig. 14), certainly
 775 when considering the improvement in the West Atlantic as described below. There
 776 is a lack of knowledge of boundary exchange processes (e.g. Arsouze et al. (2009)),
 777 which keeps open the possibility that the large sediment source from the margins is
 778 compensated by boundary exchange processes. In our model only bSiO_2 was used as
 779 a scavenger, while near the margins, other scavengers like POC and CaCO_3 might
 780 play an important role. Also scavengers not present in the PISCES model, like
 781 clay minerals (Walker et al. (1988)) might be important for reducing $[\text{Al}_{\text{diss}}]$. The
 782 possibility of this *boundary scavenging* (Bacon et al. (1988); Arsouze et al. (2009))
 783 is convincingly shown to be present in observations (e.g. Brown et al. (2010)).

784 Since most of the data used here is from the open ocean and not from coastal
 785 areas, and our model is too coarse for an analysis of the coastal aluminium con-
 786 centrations, it is not possible to make a good a priori estimate of the amount
 787 of sediment input needed for a better simulation. Since dust input and internal
 788 processes contain parameters which are, just like sediment input, not completely
 789 constrained, a change in sediment input could result in a simulation which predicts




Figure 20: Total Al budget (Tmol) in the world ocean after a partial spin-up of 600 yr of the reference simulation. From this point the reference simulation together with the sensitivity experiments is run for another 1300 yr or more. The blue solid line is the budget of the reference simulation, the red dashed line of the slow-equilibration simulation and the black dash-dotted line of the simulation with a decreased partition coefficient.

the open ocean aluminium concentration better, but does a bad job in other areas. Nevertheless, consistent with work by Moran and Moore (1991); Moran et al. (1992); Middag et al. (2011), there are clearly some areas where sediment supply of Al_{diss} appears important and a better understanding of the processes governing its supply is needed.

4.4 Internal coefficients

For the experiments where we changed two internal parameters (k_d and κ) we can now analyse the simulations in a more sophisticated manner by using the above timescale approach.

As can be seen from the black dash-dotted line in Fig. 20, the effect of a halved partition coefficient k_d on the whole ocean is an almost doubling of the total Al budget. According to Eq. 3 indeed a halved k_d means half as much $[\text{Al}_{\text{ads}}^{\text{eq}}]$. In all experiments equilibration is very fast, except the one where κ is decreased significantly. Because of the fast equilibration, with halved k_d , export will be half as fast. In a steady state, this would result in a doubled Al_{diss} content, if bSiO_2 would be distributed homogeneously. Since it is not, this does not hold for our model. At the dust deposition site in the Atlantic Ocean we see in Fig. 16(c) an increase of $[\text{Al}_{\text{diss}}]$ of less than 100% in the Atlantic Ocean near the dust deposition site. Because of the reduced k_d more Al_{diss} can be transported northward. When it reaches large $[\text{bSiO}_2]$, it is scavenged with twice the speed compared to the reference simulation. Since this extra Al_{diss} was able to reach the high $[\text{bSiO}_2]$ site, the effect of the reduced k_d is dampened and therefore the ocean Al budget is less than doubled.

If we also want to consider advection, we should look at Υ , which is proportional to k_d and $[\text{bSiO}_2]$ (see Eq. 11). The halved k_d results in only a small change in importance of scavenging relative to advection (see Fig. 19 where the logarithm of Υ is plotted).




Figure 21: $\log_{10}(\Upsilon)$, when $\kappa = 100 \text{ yr}^{-1}$ and $k_d = 4 \cdot 10^6 \text{ l/kg}$. For the advection time τ_{adv} the definition of Eq. 10 is used, and the scavenging time τ_{scav} is the maximum of τ_{trans} and τ_{sink} .

816 A decrease in the first order rate constant κ means that equilibration goes slower
817 (Eq. 2). If dust is dissolved in the surface ocean, its conversion to Al_{ads} is slower,
818 so that Al_{diss} is scavenged slower. Because of the resulting higher $[\text{Al}_{\text{diss}}]$, more
819 aluminium is transported northward by the North Atlantic Current. Fig. 6 shows
820 the biogenic silica (bSiO_2) concentration according to the model at the surface. In
821 the Atlantic Ocean, we can see large concentrations of bSiO_2 north of 40°N . The
822 dissolved Al which arrives in this area, is scavenged by this high $[\text{bSiO}_2]$. This
823 results in high concentrations at all depths around 40°N . However, because of slow
824 equilibration, Al_{diss} can go further north before it is actually scavenged. This is
825 the reason why a significant increase in $[\text{Al}_{\text{diss}}]$ is visible around 60°N at all depths,
826 and also further south at a depth around 2 km because this extra Al is advected
827 southward via the MOC (Fig. 17(d)). Also desorption goes slower, which means
828 that Al_{diss} that is adsorbed at the surface around 40°N , will desorb slowly while
829 sinking, such that compared to the reference simulation a relatively large quantity
830 of Al_{diss} will appear in the deep Atlantic Ocean. This can be seen in the lower right
831 of Fig. 17(d).

832 According to Fig. 17(d) there is a decrease of $[\text{Al}_{\text{diss}}]$ at low latitudes around 500
833 m and 1 km depth. Less dissolved Al from dust in the surface ocean is scavenged
834 and therefore less adsorbed aluminium is present in the deeper regions to desorb at
835 depth. So in the deep ocean there are regions of increased $[\text{Al}_{\text{diss}}]$ and regions of
836 decreased $[\text{Al}_{\text{diss}}]$.

837 If a lot of particulate aluminium sinks where there is almost no biogenic silica,
838 it will desorb and the resulting Al_{diss} will stay there. According to the equations,
839 with a very large κ this will happen instantaneously and the dissolved aluminium
840 will stay there indefinitely (in the limit of zero biogenic silica and zero advection),
841 i.e., $\tau_{\text{res}} = \infty$. This really means, as discussed in Section 4.2.1, that in that case
842 advection is much more important than scavenging. If κ is decreased, the particulate
843 aluminium has enough time to fall out of this low biogenic silica domain before it
844 desorbs.

845 In Fig. 21 the logarithm of the relative importance of scavenging, Υ , is plotted
846 for the slow equilibration experiment. It shows that everywhere advection is more
847 important than scavenging. Note that the residence times are based on boxes of




Figure 22: The quantities in the shaded area are (a) $[\text{Al}_{\text{diss}}]$ (nM) and (b) $[\text{Si}_{\text{diss}}]$ (μM) at the Geotraces West Atlantic cruise section. The contours represent the Atlantic Overturning Stream Function (Sv). See Fig. 9 for the large version of (a).

approximately $2^\circ \times 2^\circ \times D_{\text{ML}}$. Fig. 21 can be interpreted as that with a low κ the aluminium distribution gets more spread out, or homogenised, throughout the ocean compared with the reference experiment. As can be seen from the thick red line in Fig. 20 the relative increase of the Al budget is very small. The only strong increase is in the surface ocean. In the rest of the ocean, $[\text{Al}_{\text{diss}}]$ is just homogenised.

4.5 Al versus Si in the MOC

Our modelling results in the Atlantic Ocean show that at around 2 km depth $[\text{Al}_{\text{diss}}]$ decreases from north to south, while the concentration of dissolved silicon ($[\text{Si}_{\text{diss}}]$) increases, all the way into the Pacific Ocean, both following the Meridional Overturning Circulation (MOC). This is shown in Fig. 22, representing the concentrations of Al_{diss} and Si_{diss} in the West Atlantic Ocean. On top of these figures a contour of the Atlantic Overturning Stream Function in Sv ($1 \text{ Sv} = 10^6 \text{ m}^3\text{s}^{-1}$) of the physical forcing is plotted.

The patterns for both $[\text{Al}_{\text{diss}}]$ and $[\text{Si}_{\text{diss}}]$ are in first order easily interpreted by looking at the source of the tracer and the general currents in the Atlantic Ocean. The source of Al_{diss} is dust dissolution in the surface of the central Atlantic Ocean. It is transported northward by the Gulf Stream and the North Atlantic Current. Before the Al_{diss} reaches locations where NADW is formed, it is scavenged by biogenic silica from 40°N northward (Fig. 6) and based on our timescale analysis (Fig. 19) possibly already near 30°N . Because of remineralisation $[\text{Al}_{\text{diss}}]$ increases in the NADW. This is consistent with the fact that in the model Al_{diss} does not nicely follow the Atlantic Overturning Stream Function in the very north of the Atlantic Ocean, but rather tends to be transported into the NADW indeed near 30°N . After this the Al_{diss} is taken along with the NADW, reaching 40°S at around 2 km depth. To complete the picture of the modelled Al_{diss} distribution, low $[\text{Al}_{\text{diss}}]$ SAAMW/AAIW enters just below the surface of the Atlantic Ocean from the Southern Ocean, and low $[\text{Al}_{\text{diss}}]$ AABW into the Atlantic Ocean below 4 km depth.

The source of Si_{diss} is the large amount of diatoms that sink and remineralise south of 50°S , after which it is advected as AABW as clearly shown in Fig. 22(b). Other features visible are the SAAMW/AAIW moving below the surface of the Atlantic Ocean and low $[\text{Si}_{\text{diss}}]$ NADW diluting the dissolved Si concentration in the Atlantic Ocean. The pattern of $[\text{Si}_{\text{diss}}]$ is similar to the observations described by R. Middag et al., in preparation. The main difference is that the model overestimates $[\text{Si}_{\text{diss}}]$ in the deep North Atlantic Ocean, and is a little bit too low in the deep South Atlantic Ocean. This is consistent with the overestimated AABW inflow in the model velocity field. In reality the lower cell of the Atlantic OSF reaches a

maximum of 1 or 2 Sv, while according to our physical model forcing its maximum is about 6 Sv. Furthermore, the AABW does not go as far north as in our model. Because of this the transport of any tracer following the pathway of AABW will be too large. Therefore $[\text{Si}_{\text{diss}}]$ is overestimated in the deep North Atlantic Ocean. Coming back to the discussion in Section 3.1.1, because of the same reasons $[\text{Al}_{\text{diss}}]$ is too quickly diluted in the deep North Atlantic Ocean.

The underestimation of $[\text{Al}_{\text{diss}}]$ in the deep North Atlantic Ocean is more significant than the overestimation of $[\text{Si}_{\text{diss}}]$ in this same area. This suggests there might be another process playing a role. A likely candidate is an adjusted depth-depending sinking velocity of bSiO_2 and Al_{ads} which is depth-increasing in our current model. Recent observations (McDonnell et al. (2011)) suggest that sinking speed is not strictly depth-increasing, but we should consider other functions with depth or more sophisticated aggregation methods, as for instance done in models by Kriest and Evans (1999); Gehlen et al. (2006); Burd and Jackson (2009), but none of these efforts give results consistent with the study by McDonnell et al. (2011). Another possibility is that we need sediment sources more sophisticated than in our sensitivity experiment, as also shown by observations by Moran et al. (1992), Middag et al. (in preparation) and references therein.

5 Conclusions and outlook

The objective of this study is to come to a better understanding of the behaviour of Al, by means of simulating the Al_{diss} distribution in the ocean with a reversible scavenging and general circulation model, and comparing it to observations. In turn, these results can be used for further developing this model to simulate the Al distribution more precisely. A more realistic simulation could then be used for constraining dust deposition fields to more precisely know nutrient input rates. It can also be used to study further the influence of sediments on the water column $[\text{Al}_{\text{diss}}]$, e.g. by means of coupling the ocean model to a sediment model.

We ran the biogeochemical model PISCES off-line, forced by a climatological velocity field with a temporal resolution of five days and a monthly dust deposition field. In one of the sensitivity experiments we included a margin sediment source, which has not been done before for Al. We are able to simulate the main features of the global Al_{diss} distribution in accordance with available observations. Specifically we are able to simulate reasonably well the distribution in the West Atlantic compared to observations from the West Atlantic Geotraces cruises. Since our results are close to observations, we can assume that the Al distribution is indeed mainly controlled by advection and reversible scavenging, with bSiO_2 as the main scavenger.

It is possible to improve specific features of the distribution by changing certain parameters of the scavenging process, or adjusting the Al sources. Increasing dust Al dissolution results in an overall aluminium increase proportional to the increased dust factor everywhere in the ocean. Adding dissolution in the water column gives higher concentrations, especially near dust deposition sites around one or two km depth. Decreasing the partition coefficient k_d results in a higher concentration of Al_{diss} everywhere in the ocean. Especially the relative increase in the Al_{diss} concentration in the Southern Ocean is large. This parameter highlights the importance of the influence of spatial variability in biogenic silica on $[\text{Al}_{\text{diss}}]$, since k_d signifies the amount of Al_{diss} which can adsorb onto bSiO_2 . When we include sediments as an input to Al_{diss} , we see elevated concentrations especially near the margins, as expected. $[\text{Al}_{\text{diss}}]$ is strongly elevated in the Arctic Ocean, which is not reconcilable with the observations, and is possibly due to the lack of oceanic-margin scavengers in the model. The model should be modified such that a realistically constrained

ocean margin source of Al_{diss} is present.

By means of a timescale analysis based on our model equations, we have shown that the importance of scavenging versus advection is highly location dependent. The gridbox based residence time is strongly space dependent as well. Especially latitudinal gradients are large. This residence time varies between less than a week at high $[\text{bSiO}_2]$ and strong advection, to many years in the oligotrophic gyres.

Even though the most relevant features of the Al_{diss} distribution are captured, some fine-tuning of the key parameters is necessary. There is clearly a localised source missing in the West Atlantic between $45\text{--}50^\circ\text{N}$ near the sediment. As already hypothesised by Moran and Moore (1991), resuspension of nepheloid layers along the western boundary of the North Atlantic Ocean is a source of Al_{diss} . A more detailed analysis of the locally elevated $[\text{Al}_{\text{diss}}]$ in the West Atlantic near 45°N will be published by R. Middag et al. (in preparation).

Another location of interest is the Arctic Ocean, where we could not simulate the observations. This is possibly because we miss a sediment source, or an important process is missing in the model, namely the biological incorporation of Al into the diatom's frustules (Caschetto and Wollast (1979); Gehlen et al. (2002)). When this process is implemented in the model, dissolved Al could be taken up by diatoms in the surface ocean and released in the deep ocean when the biogenic silica is remineralised together with the incorporated Al, yielding a strong Al:Si correlation as present in the observations (Middag et al. (2009)).

We made good progress in simulating the distribution of dissolved aluminium in the world ocean, and our approach confirms that dust deposition is the main source of aluminium and reversible scavenging is the main process in removing it. However, significant improvement might be possible by (1) developing a more sophisticated model for ocean sediment source of Al, (2) adding diatom Al incorporation, (3) understanding where the elevated Al levels in the North Atlantic Ocean near the sediment come from and incorporating these sources in the model, and (4) fine tuning the free parameters and constraining them by means of field studies and laboratory experiments to better understand the role for these processes.

Acknowledgements

The authors are grateful to those who have been proved useful in discussion, among which Jack Middelburg, Laurent Bopp and Wilco Hazeleger. Furthermore, we want to thank our colleagues from the IMAU institute for their critical and useful feedback to the model configuration and output. We also want to thank the people who kindly provided their datasets that are mentioned in this paper. Christoph Heinze kindly dug up the model data from Gehlen et al. (2003) and made it available to us.

The authors wish to acknowledge use of the Ferret free software program for analysis and graphics in this paper. Ferret is a product of NOAA's Pacific Marine Environmental Laboratory (<http://ferret.pmel.noaa.gov/Ferret/>).

This research is funded by NWO, Nr. 839.08.414, to whom we are grateful.

References

- R.F. Anderson. Chemical tracers of particle transport. In H. Elderfield, editor, *The Oceans and Marine Geochemistry*, volume 6, chapter 6.09, pages 247–274. Elsevier, 2006.
- T. Arsouze, J.-C. Dutay, F. Lacan, and C. Jeandel. Reconstructing the Nd oceanic cycle using a coupled dynamical–biogeochemical model. *Biogeosciences*, 6(12): 2829–2846, 2009. ISSN 1726-4170.

- 984 O. Aumont and L. Bopp. Globalizing results from ocean in situ iron fertilization
985 studies. *Global Biogeochemical Cycles*, 20(2), 2006. ISSN 0886-6236. doi: 10.
986 1029/2005GB002591.
- 987 O. Aumont, L. Bopp, and M. Schulz. What does temporal variability in aeolian dust
988 deposition contribute to sea-surface iron and chlorophyll distributions? *Geophys.*
989 *Res. Lett.*, 35(7):L07607–, apr 2008. ISSN 00948276. doi: 10.1029/2007GL031131.
- 990 M.P. Bacon and R.F. Anderson. Distribution of thorium isotopes between dissolved
991 and particulate forms in the deep sea. *J. Geophys. Res.*, 87(C3):2045–2056, 1982.
992 doi: 10.1029/JC087iC03p02045.
- 993 M.P. Bacon, W. Roether, and H. Elderfield. Tracers of chemical scavenging in the
994 ocean: Boundary effects and large-scale chemical fractionation [and discussion].
995 *Philosophical Transactions of the Royal Society of London. Series A, Mathematical*
996 *and Physical Sciences*, 325(1583):147, 1988.
- 997 A.R. Baker, T.D. Jickells, M. Witt, and K.L. Linge. Trends in the solubility of
998 iron, aluminium, manganese and phosphorus in aerosol collected over the atlantic
999 ocean. *Marine Chemistry*, 98(1):43 – 58, 2006. ISSN 0304-4203. doi: 10.1016/j.
1000 marchem.2005.06.004.
- 1001 A.J. Van Bennekom, A.G.J. Buma, and R.F. Nolting. Dissolved aluminium in the
1002 weddell-scotia confluence and effect of al on the dissolution kinetics of biogenic
1003 silica. *Marine Chemistry*, 35(1-4):423 – 434, 1991. ISSN 0304-4203. doi: 10.1016/
1004 S0304-4203(09)90034-2.
- 1005 W.S. Broecker and T.-H. Peng. *Tracers in the Sea*. Eldigio Press, 1982.
- 1006 M.T. Brown, S.M. Lippiatt, and K.W. Bruland. Dissolved aluminum, partic-
1007 ulate aluminum, and silicic acid in northern Gulf of Alaska coastal waters:
1008 Glacial/riverine inputs and extreme reactivity. *Marine Chemistry*, 2010. doi:
1009 10.1016/j.marchem.2010.04.002.
- 1010 K.W. Bruland and M.C. Lohan. Controls of trace metals in seawater. In H. El-
1011 derfield, editor, *The Oceans and Marine Geochemistry*, volume 6, chapter 6.02,
1012 pages 23–47. Elsevier, 2006.
- 1013 A.B. Burd and G.A. Jackson. Particle aggregation. *Annual review of marine science*,
1014 1:65–90, 2009.
- 1015 S. Caschetto and R. Wollast. Vertical distribution of dissolved aluminium in the
1016 Mediterranean Sea. *Marine Chemistry*, 7(2):141–155, 1979. ISSN 0304-4203. doi:
1017 10.1016/0304-4203(79)90006-9.
- 1018 L. Chou and R. Wollast. Biogeochemical behavior and mass balance of dissolved
1019 aluminum in the western Mediterranean Sea. *Deep Sea Research Part II: Topical*
1020 *Studies in Oceanography*, 44(3-4):741 – 768, 1997. ISSN 0967-0645. doi: 10.1016/
1021 S0967-0645(96)00092-6.
- 1022 N. Dittert, L. Corrin, D. Bakker, J. Bendtsen, M. Gehlen, C. Heinze, E. Maier-
1023 Reimer, P. Michalopoulos, K.E.R. Soetaert, and R.J.S. Tol. *Integrated Data Sets*
1024 *of the EU FP5 Research Project: ORFOIS: Origin and Fate of Biogenic Particle*
1025 *Fluxes in the Ocean and Their Interactions with Atmospheric CO2 Concentra-*
1026 *tions as Well as the Marine Sediment*, volume 1. WDC-MARE Reports 0002,
1027 Alfred Wegener Institute for Polar and Marine Research, 2005.

- 1028 S. Dixit, P. Van Cappellen, and A.J. van Bennekom. Processes controlling sol-
1029 ubility of biogenic silica and pore water build-up of silicic acid in marine sed-
1030 iments. *Marine Chemistry*, 73(3-4):333–352, 2001. ISSN 0304-4203. doi:
1031 10.1016/S0304-4203(00)00118-3.
- 1032 J.C. Dutay, J.L. Bullister, S.C. Doney, J.C. Orr, R. Najjar, K. Caldeira, J.M. Campin,
1033 H. Drange, M. Follows, Y. Gao, et al. Evaluation of ocean model ventilation with
1034 cfc-11: comparison of 13 global ocean models. *Ocean Modelling*, 4(2):89–120,
1035 2002. doi: 10.1016/S1463-5003(01)00013-0.
- 1036 J.C. Dutay, F. Lacan, M. Roy-Barman, and L. Bopp. Influence of particle size
1037 and type on ^{231}Pa and ^{230}Th simulation with a global coupled biogeochemical-
1038 ocean general circulation model: A first approach. *Geochemistry Geophysics*
1039 *Geosystems*, 10(1):Q01011, 2009. doi: 10.1029/2008GC002291.
- 1040 S. Emerson and J. Hedges. Sediment diagenesis and benthic flux. In H. Elderfield,
1041 editor, *The Oceans and Marine Geochemistry*, volume 6, chapter 6.11, pages
1042 293–319. Elsevier, 2006.
- 1043 C. Ethé, O. Aumont, M-A Foujols, and M. Lévy. NEMO reference manual, tracer
1044 component : NEMO-TOP. Preliminary version. *Note du Pole de Modélisation*,
1045 *Institut Pierre-Simon Laplace*, 2006.
- 1046 M. Filella. Colloidal properties of submicron particles in natural waters. In K.J.
1047 Wilkinson and J.R. Leas, editors, *Environmental Colloids and Particles: Be-*
1048 *haviour, Separation and Characterisation*, volume 10, chapter 2, pages 17–94.
1049 John Wiley & Sons, Ltd, 2007. doi: 10.1002/9780470024539.ch2.
- 1050 Stefan F. Gary, M. Susan Lozier, Claus W. Böning, and Arne Biastoch. Deciphering
1051 the pathways for the deep limb of the Meridional Overturning Circulation. *Deep*
1052 *Sea Research Part II: Topical Studies in Oceanography*, 58(17-18):1781 – 1797,
1053 2011. ISSN 0967-0645. doi: 10.1016/j.dsr2.2010.10.059.
- 1054 M. Gehlen, L. Beck, G. Calas, A.M. Flank, A.J. van Bennekom, and J.E.E. van
1055 Beusekom. Unraveling the atomic structure of biogenic silica: evidence of the
1056 structural association of Al and Si in diatom frustules. *Geochimica et Cosmochim-*
1057 *ica Acta*, 66(9):1601–1609, 2002. doi: 10.1016/S0016-7037(01)00877-8.
- 1058 M. Gehlen, C. Heinze, E. Maier-Reimer, et al. Coupled Al-Si geochemistry in
1059 an ocean general circulation model: A tool for the validation of oceanic dust
1060 deposition fields? *Global Biogeochemical Cycles*, 17(1):1028, 2003. doi: 10.1029/
1061 2001GB001549.
- 1062 M. Gehlen, L. Bopp, N. Emprin, O. Aumont, C. Heinze, and O. Ragueneau. Rec-
1063 onciling surface ocean productivity, export fluxes and sediment composition in a
1064 global biogeochemical ocean model. *Biogeosciences*, 3(4):521–537, 2006.
- 1065 M. Gehlen, R. Gangstø, B. Schneider, L. Bopp, O. Aumont, and C. Ethe. The fate of
1066 pelagic CaCO_3 production in a high CO_2 ocean: a model study. *Biogeosciences*,
1067 4(4):505–519, 2007.
- 1068 E.D. Goldberg. Marine geochemistry 1. Chemical scavengers of the sea. *The Journal*
1069 *of Geology*, 62(3):249–265, 1954. ISSN 0022-1376.
- 1070 S. Guerzoni, E. Molinaroli, and R. Chester. Saharan dust inputs to the western
1071 Mediterranean Sea: depositional patterns, geochemistry and sedimentological im-
1072 plications. *Deep Sea Research Part II: Topical Studies in Oceanography*, 44(3-4):
1073 631–654, 1997.

- 1074 Q. Han. *Crustal Tracers in the Atmosphere and Ocean: Relating their Concentra-*
1075 *tions, Fluxes, and Ages*. PhD thesis, University of California, Irvine, 2010.
- 1076 Q. Han, J.K. Moore, C. Zender, and D. Hydes. Constraining oceanic dust deposition
1077 using surface ocean dissolved Al. *Global Biogeochemical Cycles*, 2008. doi: 10.
1078 1029/2007GB002975.
- 1079 D.J. Hydes, P.J. Statham, and J.D. Burton. A vertical profile of dissolved trace
1080 metals (Al, Cd, Cu, Mn, Ni) over the median valley of the mid Atlantic ridge,
1081 43N: Implications for Hydrothermal activity. *Science of The Total Environment*,
1082 49(0):133 – 145, 1986. ISSN 0048-9697. doi: 10.1016/0048-9697(86)90236-6.
- 1083 D.J. Hydes, G.J. de Lange, and H.J.W. de Baar. Dissolved aluminium in the
1084 Mediterranean. *Geochimica et Cosmochimica Acta*, 52(8):2107 – 2114, 1988. ISSN
1085 0016-7037. doi: 10.1016/0016-7037(88)90190-1.
- 1086 T. Jickells. Atmospheric inputs of metals and nutrients to the oceans: their mag-
1087 nitude and effects. *Marine Chemistry*, 48(3-4):199–214, 1995.
- 1088 T. Jickells, T. Church, A. Veron, and R. Arimoto. Atmospheric inputs of man-
1089 ganese and aluminium to the Sargasso Sea and their relation to surface water
1090 concentrations. *Marine chemistry*, 46(3):283–292, 1994.
- 1091 T.D. Jickells, Z.S. An, K.K. Andersen, A.R. Baker, G. Bergametti, N. Brooks, J.J.
1092 Cao, P.W. Boyd, R.A. Duce, K.A. Hunter, et al. Global iron connections between
1093 desert dust, ocean biogeochemistry, and climate. *Science*, 308(5718):67, 2005.
- 1094 J. Kramer, P. Laan, G. Sarthou, K.R. Timmermans, and H.J.W. De Baar. Distri-
1095 bution of dissolved aluminium in the high atmospheric input region of the sub-
1096 tropical waters of the North Atlantic Ocean. *Marine Chemistry*, 88(3-4):85–101,
1097 2004. ISSN 0304-4203. doi: 10.1016/j.marchem.2004.03.009.
- 1098 K. Kremling. The distribution of cadmium, copper, nickel, manganese, and alu-
1099 minium in surface waters of the open Atlantic and European shelf area. *Deep*
1100 *Sea Research Part A. Oceanographic Research Papers*, 32(5):531–555, 1985. ISSN
1101 0198-0149. doi: 10.1016/0198-0149(85)90043-3.
- 1102 I. Kriest and G.T. Evans. Representing phytoplankton aggregates in biogeochemical
1103 models. *Deep Sea Research Part I: Oceanographic Research Papers*, 46(11):1841–
1104 1859, 1999.
- 1105 P. Lam. The dynamic ocean biological pump: Insights from a global compilation
1106 of particulate organic carbon, CaCO_3 , and opal concentration profiles from the
1107 mesopelagic. *Global Biogeochem. Cycles*, 25(3):GB3009, 2011. doi: 10.1029/
1108 2010GB003868.
- 1109 J.R. Lead and K.J. Wilkinson. Environmental colloids and particles: Current
1110 knowledge and future developments. In K.J. Wilkinson and J.R. Leas, editors,
1111 *Environmental Colloids and Particles: Behaviour, Separation and Characterisa-*
1112 *tion*, volume 10, chapter 1, pages 1–16. John Wiley & Sons, Ltd, 2007. doi:
1113 10.1002/9780470024539.ch1.
- 1114 J.C. Lewin. The dissolution of silica from diatom walls. *Geochimica et Cosmochim-*
1115 *ica Acta*, 21(3-4):182 – 198, 1961. ISSN 0016-7037. doi: 10.1016/S0016-7037(61)
1116 80054-9.
- 1117 S. Loucaides, T. Behrends, and P. van Cappellen. Reactivity of biogenic silica:
1118 Surface versus bulk charge density. *Geochimica et Cosmochimica Acta*, 74(2):517
1119 – 530, 2010. ISSN 0016-7037. doi: 10.1016/j.gca.2009.10.038.

- 1120 M.S. Lozier. Deconstructing the Conveyor Belt. *Science*, 328(5985):1507, 2010. doi:
1121 10.1126/science.1189250.
- 1122 F.T. MacKenzie, M. Stoffyn, and R. Wollast. Aluminum in seawater: control by
1123 biological activity. *Science*, 199(4329):680, 1978. doi: 10.1126/science.199.4329.
1124 680.
- 1125 J.E. Mackin. Control of dissolved al distributions in marine sediments by clay recon-
1126 stitution reactions: experimental evidence leading to a unified theory. *Geochimica*
1127 *et Cosmochimica Acta*, 50(2):207–214, 1986. doi: 10.1016/0016-7037(86)90170-5.
- 1128 J.E. Mackin and R.C. Aller. The effects of clay mineral reactions on dissolved al
1129 distributions in sediments and waters of the amazon continental shelf. *Continental*
1130 *Shelf Research*, 6(1-2):245–262, 1986. doi: 10.1016/0278-4343(86)90063-4.
- 1131 G. Madec. NEMO ocean engine. *Note du Pole de Modélisation, Institut Pierre-*
1132 *Simon Laplace*, 2008.
- 1133 G. Madec, P. Delecluse, M. Imbard, and C. Lévy. OPA 8.1 Ocean General Circula-
1134 tion Model reference manual. *Note du Pole de Modélisation, Institut Pierre-Simon*
1135 *Laplace*, 11:91p, 1998.
- 1136 N. Mahowald, K. Kohfeld, M. Hansson, Y. Balkanski, S.P. Harrison, I.C. Prentice,
1137 M. Schulz, and H. Rodhe. Dust sources and deposition during the last glacial
1138 maximum and current climate: A comparison of model results with paleodata
1139 from ice cores and marine sediments. *Journal of Geophysical Research*, 104(D13):
1140 15895–15, 1999. ISSN 0148-0227.
- 1141 E. Maier-Reimer, U. Mikolajewicz, and K. Hasselmann. Mean circulation of the
1142 Hamburg LSG OGCM and its sensitivity to the thermohaline surface forcing.
1143 *Journal of Physical Oceanography*, 23(4):731–757, 1993. ISSN 0022-3670.
- 1144 H.B. Maring and R.A. Duce. The impact of atmospheric aerosols on trace metal
1145 chemistry in open ocean surface seawater, 1. aluminum. *Earth and Planetary Sci-*
1146 *ence Letters*, 84(4):381 – 392, 1987. ISSN 0012-821X. doi: 10.1016/0012-821X(87)
1147 90003-3.
- 1148 J. McAlister and K. Orians. Controls on hydroxide-speciated trace metals
1149 in the ocean, 2011. URL [http://modb.oce.ulg.ac.be/colloquium/2011/](http://modb.oce.ulg.ac.be/colloquium/2011/abstracts2011.pdf)
1150 [abstracts2011.pdf](http://modb.oce.ulg.ac.be/colloquium/2011/abstracts2011.pdf).
- 1151 A.M.P. McDonnell et al. *Marine particle dynamics: sinking velocities, size dis-*
1152 *tributions, fluxes, and microbial degradation rates*. PhD thesis, Massachusetts
1153 Institute of Technology, 2011. (no paper copy).
- 1154 C.I. Measures and Vink. On the use of dissolved aluminum in surface waters to
1155 estimate dust deposition to the ocean. *Global Biogeochem. Cycles*, 14(1):317–
1156 327, 2000. doi: 10.1029/1999GB001188.
- 1157 C.I. Measures, M.T. Brown, and S. Vink. Dust deposition to the surface waters
1158 of the western and central north pacific inferred from surface water dissolved
1159 aluminum concentrations. *Geochem. Geophys. Geosyst.*, 6(9):Q09M03–, sep 2005.
1160 ISSN 15252027. doi: 10.1029/2005GC000922.
- 1161 C.I. Measures, T. Sato, S. Vink, S. Howell, and YH Li. The fractional solubil-
1162 ity of aluminium from mineral aerosols collected in hawaii and implications for
1163 atmospheric deposition of biogeochemically important trace elements. *Marine*
1164 *Chemistry*, 120(1-4):144–153, 2010. doi: 10.1016/j.marchem.2009.01.014.

- 1165 R. Middag. *Dissolved Aluminium and Manganese in the Polar Oceans*. PhD thesis,
1166 University of Groningen, 2010.
- 1167 R. Middag, H.J.W. de Baar, P. Laan, and K. Bakker. Dissolved aluminium and
1168 the silicon cycle in the Arctic Ocean. *Marine Chemistry*, 115(3-4):176–195, 2009.
1169 doi: 10.1016/j.marchem.2009.08.002.
- 1170 R. Middag, C. van Slooten, H.J.W. de Baar, and P. Laan. Dissolved aluminium in
1171 the southern ocean. *Deep Sea Research Part II: Topical Studies in Oceanography*,
1172 2011. ISSN 0967-0645. doi: DOI:10.1016/j.dsr2.2011.03.001.
- 1173 R. Middag, H.J.W. de Baar, P. Laan, and O. Huhn. The effects of continental
1174 margins and water mass circulation on the distribution of dissolved aluminum
1175 and manganese in drake passage. *J. Geophys. Res.*, 117(C1):C01019, 2012. doi:
1176 10.1029/2011JC007434.
- 1177 J.K. Moore, O. Braucher, et al. Sedimentary and mineral dust sources of dissolved
1178 iron to the world ocean. *Biogeosciences*, 5(3):631–656, 2008.
- 1179 S.B. Moran and R. M. Moore. Temporal variations in dissolved and particulate
1180 aluminum during a spring bloom. *Estuarine, Coastal and Shelf Science*, 27(2):
1181 205 – 215, 1988. ISSN 0272-7714. doi: 10.1016/0272-7714(88)90090-X.
- 1182 S.B. Moran and R.M. Moore. The distribution of colloidal aluminum and organic
1183 carbon in coastal and open ocean waters off Nova Scotia. *Geochimica et Cos-*
1184 *mochimica Acta*, 53(10):2519–2527, 1989.
- 1185 S.B. Moran and R.M. Moore. The potential source of dissolved aluminum from re-
1186 suspended sediments to the north atlantic deep water. *Geochimica et Cosmochim-*
1187 *ica Acta*, 55(10):2745 – 2751, 1991. ISSN 0016-7037. doi: 10.1016/0016-7037(91)
1188 90441-7.
- 1189 S.B. Moran, R.M. Moore, and S. Westerlund. Dissolved aluminum in the weddell
1190 sea. *Deep Sea Research Part A. Oceanographic Research Papers*, 39(3-4):537–547,
1191 1992. doi: 10.1016/0198-0149(92)90087-A.
- 1192 H. Obata, D.S. Alibo, and Y. Nozaki. Dissolved aluminum, indium, and cerium in
1193 the Sea of Japan and the Sea of Okhotsk: Comparison to the marginal seas of
1194 the western North Pacific. *J. Geophys. Res.*, 112(C12):C12003–, dec 2007. ISSN
1195 01480227. doi: 10.1029/2006JC003944.
- 1196 K.J. Orians and K.W. Bruland. Dissolved aluminium in the central North Pacific.
1197 *Nature*, 1985. doi: 10.1038/316427a0.
- 1198 K.J. Orians and K.W. Bruland. The biogeochemistry of aluminum in the Pacific
1199 Ocean. *Earth and planetary science letters*, 78(4):397–410, 1986. ISSN 0012-821X.
1200 doi: 10.1016/0012-821X(86)90006-3.
- 1201 O. Ragueneau, P. Tréguer, A. Leynaert, R.F. Anderson, M.A. Brzezinski, D.J.
1202 DeMaster, R.C. Dugdale, J. Dymond, G. Fischer, R. Francois, et al. A review
1203 of the Si cycle in the modern ocean: recent progress and missing gaps in the
1204 application of biogenic opal as a paleoproductivity proxy. *Global and Planetary*
1205 *Change*, 26(4):317–365, 2000. ISSN 0921-8181.
- 1206 M.J.A. Rijkenberg, C.F. Powell, M. Dall’Osto, M.C. Nielsdottir, M.D. Patey, P.G.
1207 Hill, A.R. Baker, T.D. Jickells, R.M. Harrison, and E.P. Achterberg. Changes
1208 in iron speciation following a Saharan dust event in the tropical North Atlantic
1209 Ocean. *Marine Chemistry*, 110(1-2):56 – 67, 2008. ISSN 0304-4203. doi: 10.1016/
1210 j.marchem.2008.02.006.

- 1211 J.L. Sarmiento and N. Gruber. *Ocean biogeochemical dynamics*. Princeton Univer-
1212 sity Press, 2006.
- 1213 M. Schulz, A. Cozic, and S. Szopa. Lmdzt-inca dust forecast model develop-
1214 ments and associated validation efforts. In *IOP Conference Series: Earth and*
1215 *Environmental Science*, volume 7, page 012014. IOP Publishing, 2009. doi:
1216 10.1088/1755-1307/7/1/012014.
- 1217 L.O. Slemons, J.W. Murray, J. Resing, B. Paul, and P. Dutrieux. Western Pacific
1218 coastal sources of iron, manganese, and aluminum to the Equatorial Undercur-
1219 rent. *Global Biogeochemical Cycles*, 24(3), 2010. doi: 10.1029/2009GB003693.
- 1220 M. Stoffyn. Biological control of dissolved aluminum in seawater: experimental
1221 evidence. *Science*, 203(4381):651, 1979. doi: 10.1126/science.203.4381.651.
- 1222 M. Stoffyn and F.T. Mackenzie. Fate of dissolved aluminum in the oceans. *Marine*
1223 *Chemistry*, 11(2):105 – 127, 1982. ISSN 0304-4203. doi: 10.1016/0304-4203(82)
1224 90036-6.
- 1225 A. Tagliabue, L. Bopp, J.-C. Dutay, A.R. Bowie, F. Chever, P. Jean-Baptiste,
1226 E. Bucciarelli, D. Lannuzel, T. Remenyi, G. Sarthou, O. Aumont, M. Gehlen, and
1227 C. Jeandel. Hydrothermal contribution to the oceanic dissolved iron inventory.
1228 *Nature Geoscience*, 2010. doi: 10.1038/NGEO818.
- 1229 C. Textor, M. Schulz, S. Guibert, S. Kinne, Y. Balkanski, S. Bauer, T. Berntsen,
1230 T. Berglen, O. Boucher, M. Chin, et al. Analysis and quantification of the di-
1231 versities of aerosol life cycles within AeroCom. *Atmos. Chem. Phys*, 6(7):1777,
1232 2006. doi: 10.5194/acp-6-1777-2006.
- 1233 J.E.E. Van Beusekom, A.J. Van Bennekom, P. Tréguer, and J. Morvan. Aluminium
1234 and silicic acid in water and sediments of the enderby and crozet basins. *Deep*
1235 *Sea Research Part II: Topical Studies in Oceanography*, 44(5):987–1003, 1997.
1236 doi: 10.1016/S0967-0645(96)00105-1.
- 1237 S. Vink and C.I. Measures. The role of dust deposition in determining surface water
1238 distributions of Al and Fe in the South West Atlantic. *Deep Sea Research Part*
1239 *II: Topical Studies in Oceanography*, 48(13):2787 – 2809, 2001. ISSN 0967-0645.
1240 doi: 10.1016/S0967-0645(01)00018-2.
- 1241 W.J. Walker, C.S. Cronan, and H.H. Patterson. A kinetic study of aluminum
1242 adsorption by aluminosilicate clay minerals. *Geochimica et Cosmochimica Acta*,
1243 52(1):55–62, 1988.
- 1244 K.H. Wedepohl. The composition of the continental crust. *Geochimica et Cos-*
1245 *mochimica Acta*, 59(7):1217–1232, 1995. doi: 10.1016/0016-7037(95)00038-2.
- 1246 P.G. Whitehead, E.J. Wilson, and D. Butterfield. A semi-distributed integrated
1247 nitrogen model for multiple source assessment in catchments (inca): Part i model
1248 structure and process equations. *Science of The Total Environment*, 210-211(0):
1249 547 – 558, 1998. ISSN 0048-9697. doi: 10.1016/S0048-9697(98)00037-0.